

Projectively Flat Finsler Metrics of Constant Flag Curvature*

Zhongmin Shen

Revised on July 16, 2002

Abstract

Finsler metrics on an open subset in \mathbb{R}^n with straight geodesics said to be projective. It is known that the flag curvature of any projective Finsler metric is a scalar function of tangent vectors (the flag curvature must be a constant if it is Riemannian). In this paper, we discuss the classification problem on projective Finsler metrics of constant flag curvature. We express them by Taylor expansions or an algebraic formula. Many examples constructed in this paper can be used as models in Finsler geometry.

1 Introduction

The Hilbert's Fourth Problem is to characterize the (not-necessarily-reversible) distance functions on an open subset in \mathbb{R}^n such that straight lines are shortest paths [Hi]. It turns out that there are lots of solutions to the problem [Ha] [Bl] [Am] [Al] [Bu] [Po] [Sz]. The main method to characterize the solutions is the integral geometric construction using quasi-positive measures on the set of all affine hyperplanes in \mathbb{R}^n . The integral construction using symplectic forms gives another description on certain smooth solutions to the problem [Alv1][Alv2][AlFe][AlGeSm].

Distance functions induced by a Finsler metrics are regarded as *smooth* ones. Thus the Hilbert Fourth Problem in the smooth case is to characterize Finsler metrics on an open subset in \mathbb{R}^n whose geodesics are straight lines. Such Finsler metrics are called *projective Finsler metrics*. In Finsler geometry, the flag curvature $\mathbf{K}(P, \mathbf{y})$ is an analogue of the sectional curvature in Riemannian geometry. It is known that every projective Finsler metric is of scalar curvature, namely, the flag curvature $\mathbf{K}(P, \mathbf{y}) = \mathbf{K}(\mathbf{y})$ is a scalar function of tangent vectors \mathbf{y} . In the early 20th century, L. Berwald studied projective Finsler metrics of constant flag curvature, particularly in the case of zero flag curvature [Be1][Be2]. Berwald shows that the (projective) Hilbert metric on a strongly convex domain in \mathbb{R}^n has constant flag curvature $\mathbf{K} = -1$. Meanwhile P. Funk classified all projective Finsler metrics with constant flag curvature on convex domains in \mathbb{R}^2 [Fk1][Fk2]. Later on, Funk tried to show the uniqueness of projectively

*1991 Mathematics Subject Classification. 53C60, 53A20

flat Finsler metrics with $\mathbf{K} = 1$ on \mathbb{S}^2 . With additional conditions, he showed that the standard Riemannian metric is the only such metric [Fk3]. In 1995-96, R. Bryant shows that there is exactly a 2-parameter family of projectively flat Finsler metrics on \mathbb{S}^2 with $\mathbf{K} = 1$ [Br1][Br2].

Given a projective Finsler metric $F(x, y)$ on an open subset $\mathcal{U} \subset \mathbb{R}^n$, let

$$P(x, y) := \frac{F_{x^k}(x, y)y^k}{2F(x, y)}.$$

It is known that the geodesics $c(t) = (x^i(t))$ of F are characterized by

$$\frac{d^2 x^i}{dt^2} + 2P\left(x, \frac{dx}{dt}\right) \frac{dx^i}{dt} = 0.$$

See Lemma 3.1 below. We call $P(x, y)$ the *projective factor* of $F(x, y)$.

In this paper, we first give a general formula for x -analytic projective Finsler metrics $F(x, y)$ of constant flag curvature. Here $F(x, y)$ is said to be x -analytic at $x = 0$ if there is a small number $\epsilon > 0$ such that for any $y \neq 0$, $F(x, y)$ can be expressed as a power series $\sum a_{i_1 \dots i_k}(y)x^{i_1} \dots x^{i_k}$ in x . For a x -analytic projective Finsler metric $F(x, y)$ at $x = 0$, its projective factor $P(x, y)$ is x -analytic at $x = 0$ too. We are going to show that both $F(x, y)$ and $P(x, y)$ are completely determined by $F(0, y) = \psi(y)$ and $P(0, y) = \varphi(y)$.

Theorem 1.1 *Let $F(x, y)$ be a projective Finsler metric of constant flag curvature $\mathbf{K} = \lambda$ on an open neighborhood \mathcal{U} of the origin in \mathbb{R}^n . Suppose that $F(x, y)$ is x -analytic at $x = 0$. Then $F(x, y)$ and its projective factor $P(x, y)$ are given by*

$$F(x, y) = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \frac{d^m}{dt^m} [\Psi_m(y+tx)] \Big|_{t=0} \quad (1)$$

$$P(x, y) = \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \frac{d^m}{dt^m} [\Phi_m(y+tx)] \Big|_{t=0}, \quad (2)$$

where

$$\Psi_m := \begin{cases} \frac{1}{2\sqrt{-\lambda}} \left\{ (\varphi + \sqrt{-\lambda}\psi)^{m+1} - (\varphi - \sqrt{-\lambda}\psi)^{m+1} \right\} & \text{if } \lambda \neq 0 \\ (m+1)\psi\varphi^m & \text{if } \lambda = 0 \end{cases}$$

$$\Phi_m := \begin{cases} \frac{1}{2} \left\{ (\varphi + \sqrt{-\lambda}\psi)^{m+1} + (\varphi - \sqrt{-\lambda}\psi)^{m+1} \right\} & \text{if } \lambda \neq 0 \\ \varphi^{m+1} & \text{if } \lambda = 0 \end{cases}$$

where $\psi(y) := F(0, y)$ and $\varphi(y) := P(0, y)$. Conversely, for any Minkowski norm $\psi(y)$ on \mathbb{R}^n , and any positively homogeneous function of degree one $\varphi(y)$ on \mathbb{R}^n , the function $F(x, y)$ defined in (1) is a projective Finsler metric with constant flag curvature $\mathbf{K} = \lambda$ and its projective factor $P(x, y)$ is given by (2) provided that they are convergent.

In [Be2], L. Berwald proves that for any projective Finsler metric $F(x, y)$ of flag curvature $\mathbf{K} = \lambda$ and its projective factor $P(x, y)$, $F_{x^k} = (PF)_{y^k}$ and the function $f(x, y) := P(x, y) \pm \sqrt{-\lambda}F(x, y)$ satisfies

$$f_{x^k}(x, y) = f(x, y)f_{y^k}(x, y). \quad (3)$$

By these equations, we prove Theorem 1.1 in Section 4. Theorem 1.1 shows that x -analytic projective Finsler metrics of constant flag curvature $\mathbf{K} = \lambda$ are uniquely determined by $\psi(y) = F(0, y)$ and $\varphi(y) = P(0, y)$. Note that a Finsler metric $F(x, y)$ expressed in (1) is reversible if and only if $\psi(y)$ is reversible ($\psi(-y) = \psi(y)$) and $\varphi(y)$ is anti-reversible ($\varphi(-y) = -\varphi(y)$). Thus there are lots of reversible projective non-Riemannian Finsler metrics of constant flag curvature.

We can construct smooth projective Finsler metrics of constant flag curvature using algebraic equations.

Theorem 1.2 *Let $\psi(y)$ be an arbitrary Minkowski norm on \mathbb{R}^n and $\varphi(y)$ be an arbitrary positively homogeneous function of degree one on \mathbb{R}^n . Define $\Psi_{\pm}(x, y)$ by*

$$\Psi_{\pm}(x, y) = \phi_{\pm}(y + \Psi_{\pm}(x, y)x). \quad (4)$$

where $\phi_{\pm}(y) := \varphi(y) \pm \psi(y)$. Then

$$F(x, y) := \frac{1}{2} \{ \Psi_+(x, y) - \Psi_-(x, y) \} \quad (5)$$

is a projective Finsler metric with constant flag curvature $\mathbf{K} = -1$ and $F(0, y) = \psi(y)$. Its projective factor $P(x, y)$ is given by

$$P(x, y) = \frac{1}{2} \{ \Psi_+(x, y) + \Psi_-(x, y) \} \quad (6)$$

with $P(0, y) = \varphi(y)$.

The proof of Theorem 1.2 is given in Section 5. We first prove that the functions $\Psi_{\pm}(x, y)$ defined in (4) satisfy (3), i.e., $(\Psi_{\pm})_{x^k} = \Psi_{\pm}(\Psi_{\pm})_{y^k}$, then using this equation, we show that the functions $F(x, y)$ and $P(x, y)$ defined in (5) and (6) have the desired properties. That is, we construct a projective Finsler metrics of constant flag curvature $\mathbf{K} = -1$ using an algebraic equation (4) for any given data $\{\varphi(y), \psi(y)\}$. By the formulas in Theorem 1.2, we obtain many interesting examples.

Theorem 1.3 *Let $\psi(y)$ be an arbitrary Minkowski norm on \mathbb{R}^n and $\varphi(y)$ be an arbitrary positively homogeneous function of degree one on \mathbb{R}^n . Define $P(x, y)$ by*

$$P(x, y) = \varphi(y + P(x, y)x). \quad (7)$$

Let

$$F(x, y) := \psi\left(y + P(x, y) x\right) \left\{1 + P_{y^m}(x, y)x^m\right\}. \quad (8)$$

Then $F(x, y)$ is a projective Finsler metric zero flag curvature and $F(0, y) = \psi(y)$ and its projective factor is $P(x, y)$ with $P(0, y) = \varphi(y)$.

The proof of Theorem 1.3 is given in Section 6. In [Be2], L. Berwald shows that any projective Finsler metric $F(x, y)$ of zero flag curvature and its projective factor, $P(x, y)$, satisfy $F_{x^k} = (PF)_{y^k}$ and $P_{x^k} = PP_{y^k}$. Then Berwald gives a formula in certain special cases. In particular, he constructs a nontrivial projective Finsler metric of zero flag curvature. See (49) below. In Theorem 1.3, we construct a projective Finsler metric of zero flag curvature for any given data $\{\varphi(y), \psi(y)\}$. By the formulas in Theorem 1.3, we obtain many interesting examples.

The construction of projective Finsler metrics of positive constant curvature is much more complicated. We will discuss this issue in Section 7.

The above theorems only characterize the local metric structure of projective Finsler metrics of constant flag curvature. When we impose some extra global conditions, the projective Finsler metrics of constant curvature are much more special. For example, H. Akbar-Zadeh proves that any Finsler metric of negative constant flag curvature on a *compact* manifold must be Riemannian and any Finsler metric of zero curvature on a *compact* manifold is locally Minkowskian [AZ].

The Beltrami theorem says that a Riemannian metric is locally projectively flat if and only if it is of constant flag curvature. For Finsler metrics, this is no longer true. The first family of non-projectively flat Finsler metrics of constant flag curvature are constructed in [BaSh]. Later on, the author constructs infinitely many non-projectively flat Finsler metrics with constant flag curvature [Sh2][Sh3]. So far, many known non-projectively flat Finsler metrics of constant flag curvature are in the form $F = \alpha + \beta$, where α is a Riemannian metric and β is a 1-form. Such metrics are called *Randers metrics* [Ran]. Recently, D. Bao and C. Robles [BaRo] have found an equivalent condition for Randers metrics be of constant flag curvature. See also [MaSh]. However, the classification problem for Randers metrics of constant flag curvature has not been completely solved yet.

After the preliminary version of this paper was sent out in the summer of 2001, the author received an interesting paper from B. Bryant [Br3], in which Bryant characterizes the (generalized) Finsler metrics on S^n with $\mathbf{K} = 1$ and great circles being as geodesics (see Theorem 2 in [Br3]). As argued by Bryant, the idea of Theorem 2 in [Br3] can be used to construct projective Finsler metrics with constant flag curvature $\mathbf{K} = 1$ and a prescribed tangent indicatrix at one point (see Proposition 4 in [Br3]). He also shows how to construct all possible local projective Finsler metrics with $\mathbf{K} = 1$. In a private communication,

Bryant informed the author that his idea can also be used to characterize (local) projective Finsler metrics of constant flag curvature $\mathbf{K} = 0$ or -1 , although he did not give any detailed discussion on this issue in [Br3]. Nevertheless, our results overlap Bryant's results in the positive flag curvature case. Here we only discuss the local construction, while Bryant in [Br3] discusses both the local and global issues. We should point out that our approach is different from Bryant's one, hence the expressions of our examples are also different his.

Many Finsler metrics with special curvature properties exist only locally. For example, there are lots of reversible projective Finsler metrics with $\mathbf{K} = 1$ on an open subset of S^n , which can not be extended to the whole S^n . R. Bryant shows that the only reversible projective Finsler metric with $\mathbf{K} = 1$ on S^2 is the standard Riemannian metric [Br1][Br2]. Thank the referee for pointing out that the above problem in higher dimensions can be resolved by the discussion around Bryant's Example 2 in [Br3]. Namely, the only reversible projective Finsler metric with $\mathbf{K} = 1$ on S^n is the standard Riemannian metric.

Acknowledgments: This work was done during author's visit to Prof. S.S. Chern. It was presented at the 2nd national conference on Finsler geometry in Nankai Institute of Mathematics on August 1, 2001. The author would like to thank S.S. Chern for his warm hospitality and valuable discussion. Thank R. Bryant and Z.I. Szabo for explaining their work to the author.

2 Preliminaries

A Minkowski norm $\psi(y)$ on a vector space V is a C^∞ function on $V \setminus \{0\}$ with the following properties: (a) $\psi(y) \geq 0$ and $\psi(y) = 0$ if and only if $y = 0$; (b) $\psi(y)$ is positively homogeneous function of degree one, i.e., $\psi(ty) = t\psi(y)$, $t \geq 0$; (c) $\psi(y)$ is strongly convex, i.e., for any $y \neq 0$, the matrix $g_{ij}(y) := \frac{1}{2}[F^2]_{y^i y^j}(y)$ is positive definite. A Finsler metric F on a manifold M is C^∞ function on $TM \setminus \{0\}$ such that $F_x := F|_{T_x M}$ is a Minkowski norm on $T_x M$ for any $x \in M$.

Let F be a Finsler metric on an n -dimensional manifold M . For a non-zero vector $\mathbf{y} \in T_p M$, F induces an inner product $g_{\mathbf{y}}$ on $T_p M$ by

$$g_{\mathbf{y}}(\mathbf{u}, \mathbf{v}) := g_{ij}(x, y)u^i v^j = \frac{1}{2}[F^2]_{y^i y^j}(x, y)u^i v^j.$$

Here $x = (x^i)$ denotes the coordinates of $p \in M$ and $(x, y) = (x^i, y^i)$ denotes the local coordinates of $\mathbf{y} \in T_p M$. The geodesics are characterized by

$$\frac{d^2 x^i}{dt^2} + 2G^i\left(x(t), \frac{dx}{dt}(t)\right) = 0,$$

where $G^i := \frac{1}{2}g^{il}\{[F^2]_{x^k y^l}y^k - [F^2]_{x^l}\}$ are called the *geodesic coefficients* of F . F is said to be *positively complete* (resp. *complete*) if every geodesic defined on (a, b) can be extended to a geodesic defined on (a, ∞) (resp. $(-\infty, \infty)$). Many examples found in this paper are positively complete, but not complete.

The Riemann curvature $\mathbf{R}_y = R^i_k dx^k \otimes \frac{\partial}{\partial x^i}|_p : T_p M \rightarrow T_p M$ is a family of linear maps on tangent spaces, defined by

$$R^i_k = 2 \frac{\partial G^i}{\partial x^k} - y^j \frac{\partial^2 G^i}{\partial x^j \partial y^k} + 2G^j \frac{\partial^2 G^i}{\partial y^j \partial y^k} - \frac{\partial G^i}{\partial y^j} \frac{\partial G^j}{\partial y^k}. \quad (9)$$

For a tangent plane $P \subset T_p M$ and a non-zero vector $\mathbf{y} \in T_p M$, the *flag curvature* $\mathbf{K}(P, y)$ is defined by

$$\mathbf{K}(P, \mathbf{y}) := \frac{g_{\mathbf{y}}(\mathbf{u}, \mathbf{R}_{\mathbf{y}}(\mathbf{u}))}{g_{\mathbf{y}}(\mathbf{y}, \mathbf{y})g_{\mathbf{y}}(\mathbf{u}, \mathbf{u}) - g_{\mathbf{y}}(\mathbf{y}, \mathbf{u})^2}, \quad (10)$$

where $P = \text{span}\{\mathbf{y}, \mathbf{u}\}$. When F is Riemannian, $\mathbf{K}(P, \mathbf{y}) = \mathbf{K}(P)$ is independent of $\mathbf{y} \in P$. It is just the sectional curvature of P in Riemannian geometry. We say that F is of *scalar curvature* if for any $\mathbf{y} \in T_p M$, the flag curvature $\mathbf{K}(P, \mathbf{y}) = \mathbf{K}(\mathbf{y})$ is independent of P containing $\mathbf{y} \in T_p M$, that is equivalent to the following system of equations in a local coordinate system (x^i, y^i) in TM ,

$$R^i_k = \mathbf{K} F^2 \left\{ \delta_k^i - F^{-1} F_{y^k} y^i \right\}.$$

If \mathbf{K} is a constant, then F is said to be of *constant flag curvature*.

There are several non-Riemannian quantities in Finsler geometry. One of the important non-Riemannian quantities is the E-curvature $\mathbf{E}_y = E_{ij} dx^i \otimes dx^j|_p : T_p M \otimes T_p M \rightarrow \mathbb{R}$, defined by

$$E_{ij} := \frac{1}{2} \frac{\partial^2}{\partial y^i \partial y^j} \left[\frac{\partial G^m}{\partial y^m} \right]. \quad (11)$$

The E-curvature is closely related to the flag curvature. For a two-dimensional plane $P \subset T_p M$ and a non-zero vector $\mathbf{y} \in T_p M$, the *flag E-curvature* $\mathbf{E}(P, y)$ is defined by

$$\mathbf{E}(P, \mathbf{y}) := \frac{F^3(\mathbf{y}) \mathbf{E}_{\mathbf{y}}(\mathbf{u}, \mathbf{u})}{g_{\mathbf{y}}(\mathbf{y}, \mathbf{y})g_{\mathbf{y}}(\mathbf{u}, \mathbf{u}) - g_{\mathbf{y}}(\mathbf{y}, \mathbf{u})^2}, \quad (12)$$

where $P = \text{span}\{\mathbf{y}, \mathbf{u}\}$. We say that F has constant flag E-curvature if for any flag (P, \mathbf{y}) , $\mathbf{E}(P, \mathbf{y}) = (n+1)c$, that is equivalent to the following system of equations,

$$E_{ij} = (n+1)c F_{y^i y^j}.$$

We know that the Funk metric on a strongly convex domain satisfies that $\mathbf{K} = -1/4$ and $\mathbf{E} = (n+1)/4$ [Sh1] (see Example 5.3 below). There are many Finsler manifolds of constant flag curvature and constant flag E-curvature [Sh2][Sh3].

3 Projective metrics

A distance function d on a manifold M is said to be *smooth* if it is induced by a Finsler metric F on M ,

$$d(p, q) := \inf_c \int_0^1 F(\dot{c}(t)) dt,$$

where the infimum is taken over all curves $c(t)$, $0 \leq t \leq 1$, joining $p = c(0)$ to $q = c(1)$. Thus smooth distance functions can be studied using calculus (see [AIM], [BCS]). The Hilbert Fourth Problem in the smooth case is to characterize Finsler metrics on an open subset in \mathbb{R}^n whose geodesics are straight lines.

Let $\mathcal{U} \subset \mathbb{R}^n$ be an open subset. A Finsler metric on \mathcal{U} is a function on $T\mathcal{U} \cong \mathcal{U} \times \mathbb{R}^n$. It is known that a Finsler metric $F(x, y)$ on \mathcal{U} is projective if and only if its geodesic coefficients G^i are in the form

$$G^i(x, y) = P(x, y)y^i,$$

where $P(x, y)$ is positively y -homogeneous with degree one, $P(x, ky) = kP(x, y)$, $k > 0$. We call $P(x, y)$ the *projective factor* of $F(x, y)$. The following lemma plays an important role.

Lemma 3.1 ([Ha][Rap]) *Let $F(x, y)$ be a Finsler metric on an open subset $\mathcal{U} \subset \mathbb{R}^n$. $F(x, y)$ is projective on \mathcal{U} if and only if it satisfies*

$$F_{x^k y^i}(x, y)y^k = F_{x^i}(x, y). \quad (13)$$

In this case, the projective factor $P(x, y)$ is given by

$$P(x, y) = \frac{F_{x^k}(x, y)y^k}{2F(x, y)}. \quad (14)$$

Let $F(x, y)$ be a projective Finsler metric on an open subset $\mathcal{U} \subset \mathbb{R}^n$ and $P(x, y)$ denote its projective factor. Plugging $G^i = Py^i$ into (9) yields

$$R^i_k = \Xi \delta_k^i + \tau_k y^i, \quad (15)$$

where

$$\Xi := P^2 - P_{x^k} y^k, \quad \tau_k := 3(P_{x^k} - PP_{y^k}) + \Xi_{y^k}. \quad (16)$$

By the symmetry property that $g_{ji}R^i_k = g_{ki}R^i_j$, we obtain

$$R^i_k = \Xi \left\{ \delta_k^i - F^{-1} F_{y^k} y^i \right\}. \quad (17)$$

Thus

$$P_{x^k} - PP_{y^k} = -\frac{(\Xi F)_{y^k}}{3F}. \quad (18)$$

From (17), we see that F is of scalar curvature with flag curvature

$$\mathbf{K} = \Xi/F^2. \quad (19)$$

This observation is made by L. Berwald [Be2]. The following lemma follows from his arguments.

Lemma 3.2 *Let $F(x, y)$ be a Finsler metric on an open subset $\mathcal{U} \subset \mathbb{R}^n$. Then $F = F(x, y)$ is projective with flag curvature $\mathbf{K} = \mathbf{K}(x, y)$ if and only if there is a positively y -homogeneous function with degree one, $P = P(x, y)$, and a positively homogeneous function of degree zero, $\mathbf{K} = \mathbf{K}(x, y)$, on $T\mathcal{U} \cong \mathcal{U} \times \mathbb{R}^n$ such that*

$$F_{x^k} = (PF)_{y^k} \quad (20)$$

$$P_{x^k} = PP_{y^k} - \frac{1}{3F}(\mathbf{K}F^3)_{y^k}. \quad (21)$$

In this case, P is the projective factor of F .

4 Analytic solutions

In this section, we are going to determine the local structures of x -analytic projective Finsler metrics $F(x, y)$ of constant flag curvature by solving the first order partial differential equations (20) and (21).

Let $F(x, y)$ be a projective Finsler metric of constant flag curvature $\mathbf{K} = \lambda$ on a neighborhood of the origin in \mathbb{R}^n and $P(x, y)$ be the projector factor. It follows from Lemma 3.2 that

$$F_{x^k} = (PF)_{y^k}, \quad P_{x^k} = PP_{y^k} - \lambda FF_{y^k}. \quad (22)$$

Let $\Psi_m(y)$ and $\Phi_m(y)$ be defined in Theorem 1.1.

Assume that $\lambda \neq 0$. Let $\Psi_{\pm}(x, y) := P(x, y) \pm \sqrt{-\lambda}F(x, y)$. Then we can express $F(x, y)$ and $P(x, y)$ in terms of $\Psi_{\pm}(x, y)$. It follows from (22) that

$$(\Psi_{\pm})_{x^k} = \Psi_{\pm}(\Psi_{\pm})_{y^k}. \quad (23)$$

The above observation is due to Berwald [Be2].

Now we are going to use (22) and (23) to find the Taylor expansion for x -analytic projective Finsler metrics of constant flag curvature $\mathbf{K} = \lambda \neq 0$.

Differentiating $\Psi_{\pm}(x, y)$ and using (23), we obtain

$$(\Psi_{\pm})_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{m+1} \left[(\varphi \pm \sqrt{-\lambda} \psi)^{m+1} \right]_{y^{i_1} \dots y^{i_m}}(y). \quad (24)$$

By (24) we obtain

$$F_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{(m+1)} \left[\Psi_m \right]_{y^{i_1} \dots y^{i_m}}(y) \quad (25)$$

$$P_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{(m+1)} \left[\Phi_m \right]_{y^{i_1} \dots y^{i_m}}(y). \quad (26)$$

By (25) and (26), we immediately obtain (1) and (2).

Now assume that $\lambda = 0$. It follows from (22) that

$$F_{x^k} = (PF)_{y^k}, \quad P_{x^k} = PP_{y^k}. \quad (27)$$

By the first equation of (27), we have

$$F_{x^{i_1} \dots x^{i_m}}(x, y) = (FP^m)_{y^{i_1} \dots y^{i_m}}(x, y). \quad (28)$$

Setting $x = 0$ in (28) yields

$$F_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{(m+1)} [\Psi_m]_{y^{i_1} \dots y^{i_m}}(y).$$

Since $P(x, y)$ satisfies the second equation of (27) with $P(0, y) = \varphi(y)$, we have

$$P_{x^{i_1} \dots x^{i_m}}(0, y) = \frac{1}{m+1} [\Phi_m]_{y^{i_1} \dots y^{i_m}}(y).$$

By the above identities, we obtain (1) and (2) in the case when $\lambda = 0$.

The second part of the theorem is left for the reader to verify. Q.E.D.

5 $\mathbf{K} = -1$

In this section, we are going to construct smooth solutions $F(x, y)$ of constant flag curvature $\mathbf{K} = -1$ with $F(0, y) = \psi(y)$ and $F_{x^k}(0, y)y^k = 2\psi(y)\varphi(y)$ for any given pair $\{\psi(y), \varphi(y)\}$. We need the following lemma.

Lemma 5.1 *Let $\phi(y)$ be an arbitrary positively homogeneous function of degree one on \mathbb{R}^n . Suppose that ϕ is C^∞ on $\mathbb{R}^n \setminus \{0\}$. There is a unique real-valued function $f := f(x, y)$ satisfying the following*

$$f(x, y) = \phi(y + f(x, y)x). \quad (29)$$

Moreover, f satisfies

$$f_{x^k} = f f_{y^k}. \quad (30)$$

Proof: Let $h(t) := t - \phi(y + tx)$. Observe that there is a small $\delta > 0$ such that for any $x \in \mathbb{R}^n$ with $|x| < \delta$, at any t where $y + tx \neq 0$,

$$h'(t) = 1 - \phi_{y^i}(y + tx)x^i \geq \frac{1}{2}.$$

Thus there is a unique t_o such that $h(t_o) = 0$. Setting $f(x, y) := t_o$ we obtain the unique solution.

Differentiating (29) with respect to x^k and y^k , respectively, we obtain

$$\begin{aligned} (1 - \phi_{y^m} x^m) f_{x^k} &= \phi_{y^k} f \\ (1 - \phi_{y^m} x^m) f_{y^k} &= \phi_{y^k}. \end{aligned}$$

For x close to 0, $\phi_{y^m} x^m < 1$. Thus (30) holds.

Q.E.D.

Proof of Theorem 1.2: Let $\Psi_{\pm}(x, y)$ be the functions defined in (4). By Lemma 5.1, $\Psi_{\pm} = \Psi_{\pm}(x, y)$ satisfy (30), i.e.,

$$(\Psi_{\pm})_{x^k} = \Psi_{\pm}(\Psi_{\pm})_{y^k} \quad (31)$$

with $\Psi_{\pm}(0, y) = \phi_{\pm}(y) = \varphi(y) \pm \psi(y)$. Let $F := \frac{1}{2}\{\Psi_+ - \Psi_-\}$ be defined in (5). It follows from (31) that

$$(\Psi_{\pm})_{x^k y^l} y^k = (\Psi_{\pm})_{x^l}.$$

Thus $F(x, y)$ satisfies (13) and it is projective by Lemma 3.1. Observe that

$$F_{x^k} y^k = \frac{1}{2} \left\{ \Psi_+(\Psi_+)_{y^k} - \Psi_-(\Psi_-)_{y^k} \right\} y^k = \frac{1}{2} \left\{ \Psi_+^2 - \Psi_-^2 \right\}.$$

Thus the projective factor $P = \frac{1}{2} F^{-1} F_{x^k} y^k$ is given by

$$P := \frac{F_{x^k} y^k}{2F} = \frac{1}{2} \frac{\Psi_+^2 - \Psi_-^2}{\Psi_+ - \Psi_-} = \frac{1}{2} \left\{ \Psi_+ + \Psi_- \right\}.$$

By a similar argument, we obtain

$$\Xi = P^2 - P_{x^k} y^k = - \left(\frac{\Psi_+ - \Psi_-}{2} \right)^2 = -F^2.$$

Thus the flag curvature $\mathbf{K} = -1$ by (19).

Q.E.D.

Example 5.1 Let $\phi(y)$ be a Minkowski norm on \mathbb{R}^n and $\mathcal{U} := \left\{ y \in \mathbb{R}^n \mid \phi(y) < 1 \right\}$. Define $\Theta(x, y)$ by

$$\Theta(x, y) = \phi \left(y + \Theta(x, y) x \right), \quad y \in T_x \mathcal{U}. \quad (32)$$

$\Theta(x, y)$ is a Finsler metric on \mathcal{U} satisfying the following system of equations [Ok],

$$\Theta_{x^k} = \Theta \Theta_{y^k}. \quad (33)$$

By Lemma 3.2, $F(x, y) := \Theta(x, y)$ is a projective Finsler metric on \mathcal{U} with constant flag curvature $\mathbf{K} = -1/4$ and its projective factor is given by $P(x, y) := \frac{1}{2} \Theta(x, y)$.

Let $\phi_+ = \phi$ and ϕ_- be an arbitrary positively y-homogeneous function of degree one on \mathbb{R}^n . Let

$$\varphi(y) := \frac{1}{2} \left\{ \phi_+(y) + \phi_-(y) \right\}, \quad \psi(y) := \frac{1}{2} \left\{ \phi_+(y) - \phi_-(y) \right\}.$$

Let $\Psi_{\pm}(x, y)$ be the solutions of (4). Let $F(x, y)$ and $P(x, y)$ be the functions defined in (5) and (6) using $\Psi_{\pm}(x, y)$. By Theorem 1.2, $F(x, y)$ is projective with $\mathbf{K} = -1$ and its projective factor is $P(x, y)$.

Note that $\Psi_+(x, y) = \Theta(x, y)$. If $\phi_-(y) := -\phi(-y)$, then $\Psi_-(x, y) = -\Theta(x, -y)$. In this case,

$$F(x, y) = \frac{1}{2} \left\{ \Theta(x, y) + \Theta(x, -y) \right\} \quad (34)$$

$$P(x, y) = \frac{1}{2} \left\{ \Theta(x, y) - \Theta(x, -y) \right\}. \quad (35)$$

We call F the *Hilbert metric* on \mathcal{U} .

Example 5.2 Let $\phi(y)$ be an arbitrary Minkowski norm on \mathbb{R}^n and $\Theta(x, y)$ denote the Funk metric of ϕ defined by (32). Let $\delta \neq 0, \lambda$ be constants and

$$\psi(y) := \frac{1}{2} \left\{ \phi(y) - \delta\phi(\lambda y) \right\}, \quad \varphi(y) := \frac{1}{2} \left\{ \phi(y) + \delta\phi(\lambda y) \right\}.$$

We assume that δ and λ are such that $\psi(y)$ is a Minkowski norm on \mathbb{R}^n . We have

$$(\varphi + \psi)(y) = \phi(y), \quad (\varphi - \psi)(y) = \delta\phi(\lambda y).$$

Let $\Psi_+(x, y)$ and $\Psi_-(x, y)$ be defined in (4) with $\phi_+(y) = \phi(y)$ and $\phi_-(y) := \delta\phi(\lambda y)$, respectively. Then

$$\Psi_+(x, y) = \Theta(x, y), \quad \Psi_-(x, y) = \delta\Theta(\delta\lambda x, \lambda y).$$

By Theorem 1.2, we conclude that the following function

$$F(x, y) := \frac{1}{2} \left\{ \Theta(x, y) - \delta\Theta(\delta\lambda x, \lambda y) \right\} \quad (36)$$

is a projective Finsler metric on its domain with $\mathbf{K} = -1$ and its projective factor is given by

$$P(x, y) = \frac{1}{2} \left\{ \Theta(x, y) + \delta\Theta(\delta\lambda x, \lambda y) \right\}. \quad (37)$$

By taking $\delta = -1$ and $\lambda = -1$ in (36), we obtain the Hilbert metric.

Take a look at the special case when $\phi(y) = |y|$, the Funk metric on \mathbb{B}^n is given by

$$\Theta(x, y) = \frac{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2}, \quad y \in T_x\mathbb{B}^n. \quad (38)$$

Let $\epsilon := \delta|\lambda|$. Then the Finsler metric defined in (36) is given by

$$F(x, y) = \frac{1}{2} \left\{ \frac{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2} - \frac{\epsilon\sqrt{|y|^2 - \epsilon^2(|x|^2|y|^2 - \langle x, y \rangle^2)} + \epsilon^2\langle x, y \rangle}{1 - \epsilon^2|x|^2} \right\}. \quad (39)$$

Clearly, F is positively complete on $\mathbb{B}^n(1)$ if $|\epsilon| < 1$ and F is the Klein metric when $\epsilon = -1$. By the above argument, we know that F is projective with constant flag curvature $\mathbf{K} = -1$.

Example 5.3 Let $\phi(y)$ be an arbitrary Minkowski norm on \mathbb{R}^n and $\Theta(x, y)$ denote the Funk metric of ϕ . For a constant vector $a \in \mathbb{R}^n$, let

$$\psi(y) := \frac{1}{2}(\phi(y) + \langle a, y \rangle), \quad \varphi(y) := \frac{1}{2}(\phi(y) - \langle a, y \rangle),$$

such that

$$\varphi(y) + \psi(y) = \phi(y), \quad \varphi(y) - \psi(y) = -\langle a, y \rangle.$$

Let $\Psi_+(x, y)$ and $\Psi_-(x, y)$ be the function defined by (4) with $\phi_+(y) = \phi(y)$ and $\phi_-(y) = -\langle a, y \rangle$, respectively. We have

$$\Psi_+(x, y) = \Theta(x, y), \quad \Psi_-(x, y) = -\frac{\langle a, y \rangle}{1 + \langle a, x \rangle}.$$

By Theorem 1.2, we know that the following function

$$F(x, y) = \frac{1}{2} \left\{ \Theta(x, y) + \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right\} \quad (40)$$

is a projective Finsler metric with $\mathbf{K} = -1$ and its projective factor is given by

$$P(x, y) = \frac{1}{2} \left\{ \Theta(x, y) - \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right\}. \quad (41)$$

Since the geodesic coefficients are in the form $G^i = Py^i$, a direct computation gives

$$\frac{\partial G^m}{\partial y^m} = (n+1)P.$$

Thus

$$E_{ij} = \frac{1}{2} \frac{\partial^2}{\partial y^i \partial y^j} \left[\frac{\partial G^m}{\partial y^m} \right] = \frac{n+1}{2} P_{y^i y^j}.$$

By (33) again, one immediately obtains

$$P(x, y) = \frac{1}{2} \left\{ \Theta(x, y) - \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right\}.$$

Observe that

$$\begin{aligned} E_{ij} &= \frac{n+1}{4} \left[\Theta - \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right]_{y^i y^j} \\ &= \frac{n+1}{4} \left[\Theta + \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right]_{y^i y^j} \\ &= \frac{n+1}{2} F_{y^i y^j}. \end{aligned}$$

Thus the Finsler metric F in (40) also has constant flag E-curvature, $\mathbf{E} = \frac{1}{2}(n+1)$.

Take a look at the special case when $\phi(y) = |y|$. The Funk metric $\Theta(x, y)$ on the unit ball $\mathbb{B}^n(1)$ is given by (38). Thus for any constant vector $a \in \mathbb{R}^n$ with $|a| < 1$, the following function

$$F(x, y) = \frac{1}{2} \left\{ \frac{\sqrt{|y|^2 - (|x|^2|y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2} + \frac{\langle a, y \rangle}{1 + \langle a, x \rangle} \right\} \quad (42)$$

is a projective Finsler metric on $\mathbb{B}^n(1)$ with $\mathbf{K} = -1$ and $\mathbf{E} = \frac{1}{2}(n+1)$. In [Sh4], we prove that a Randers metric is projective with constant flag curvature if and only if it is locally Minkowskian or, up to a scaling, isometric to the metric in (42).

6 $\mathbf{K} = 0$

In this section, we are going to prove Theorem 1.3 then construct some interesting projective Finsler metric $F(x, y)$ with zero flag curvature $\mathbf{K} = 0$.

Proof of Theorem 1.3: Let $P(x, y)$ and $F(x, y)$ be defined in (7) and (8) respectively. By Lemma 5.1, $P(x, y)$ satisfies

$$P_{x^k} = PP_{y^k} \quad (43)$$

with $P(0, y) = \varphi(y)$. It follows from (43) that

$$P_{x^k y^m} = \frac{1}{2}(P^2)_{y^k y^m} = P_{x^m y^k}. \quad (44)$$

Differentiating (8) with respect to x^k and using (43), we obtain

$$\begin{aligned} F_{x^k} &= \psi_{y^m}(y + Px) \left\{ P\delta_k^m + P_{x^k} x^m \right\} \left\{ 1 + P_{y^l} x^l \right\} \\ &\quad + \psi(y + Px) \left\{ P_{y^k} + P_{y^m x^k} x^m \right\}. \end{aligned}$$

It follows from (43) and (8) that

$$PF = \psi(y + Px) \left\{ P + P_{x^m} x^m \right\}. \quad (45)$$

Differentiating (45) with respect to y^k yields

$$\begin{aligned} (PF)_{y^k} &= \psi_{y^m}(y + Px) \left\{ \delta_k^m + P_{y^k} x^m \right\} \left\{ P + P_{x^l} x^l \right\} \\ &\quad + \psi(y + Px) \left\{ P_{y^k} + P_{x^m y^k} x^m \right\}. \end{aligned}$$

By (43), we have

$$\left\{ P\delta_k^m + P_{x^k} x^m \right\} \left\{ 1 + P_{y^l} x^l \right\} = \left\{ \delta_k^m + P_{y^k} x^m \right\} \left\{ P + P_{x^l} x^l \right\}.$$

In virtue of (44), we conclude that F satisfies

$$F_{x^k} = (PF)_{y^k}. \quad (46)$$

By Lemma 3.2, we conclude that the function $F(x, y)$ is a projective Finsler metric on its domain with $\mathbf{K} = 0$ and its projective factor is $P(x, y)$. Q.E.D.

Example 6.1 Let $\phi(y)$ be a Minkowski norm on \mathbb{R}^n and $\Theta(x, y)$ denote the Funk metric on $\mathcal{U} := \{y \in \mathbb{R}^n \mid \phi(y) < 1\}$ which is defined in (32). Let $\psi(y) = \phi(y) + \langle a, y \rangle$ and $\varphi(y) = \phi(y)$. Let $P(x, y)$ and $F(x, y)$ be defined in (7) and (8) respectively. We have

$$P(x, y) = \Theta(x, y).$$

Observe that

$$\phi(y + Px) = \phi(y + \Theta x) = \Theta.$$

Thus

$$\psi(y + Px) = \Theta + \langle a, y \rangle + \langle a, x \rangle \Theta.$$

This gives

$$F(x, y) = \left\{ \left(1 + \langle a, x \rangle \right) \Theta(x, y) + \langle a, y \rangle \right\} \left\{ 1 + \Theta_{y^k}(x, y) x^k \right\}.$$

Take $\psi(y) = |y| + \langle a, y \rangle$ and $\varphi(y) = |y|$. We obtain

$$\begin{aligned} F(x, y) &= \left\{ 1 + \langle a, x \rangle + \frac{(1 - |x|^2) \langle a, y \rangle}{\sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle} \right\} \\ &\quad \times \frac{\left(\sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle \right)^2}{(1 - |x|^2)^2 \sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)}}. \end{aligned} \quad (47)$$

By Theorem 1.3, we know that $F(x, y)$ is a projective Finsler metric with zero flag curvature $\mathbf{K} = 0$ and its projective factor is given by

$$P(x, y) = \frac{\sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle}{1 - |x|^2}. \quad (48)$$

Clearly, $F(x, y)$ is not locally Minkowskian.

Remark 6.1 Letting $a = 0$ in (47), we obtain

$$F(x, y) = \frac{\left(\sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)} + \langle x, y \rangle \right)^2}{(1 - |x|^2)^2 \sqrt{|y|^2 - (|x|^2 |y|^2 - \langle x, y \rangle^2)}}. \quad (49)$$

This is just the projective Finsler metric constructed by L. Berwald [Be2]. The Finsler metric $F(x, y)$ in (49) is positively complete, i.e., every unit speed geodesic on (δ, τ) can be extended to a geodesic on (δ, ∞) . It can be easily shown that any complete projective Finsler metric with zero flag curvature $\mathbf{K} = 0$ must be a Minkowski metric [Be1][Fk1].

Example 6.2 Let ϕ be an arbitrary Minkowski norm on \mathbb{R}^n . Take

$$\psi(y) := \phi(y), \quad \varphi(y) := -\langle a, y \rangle.$$

We obtain

$$P(x, y) = -\frac{\langle a, y \rangle}{1 + \langle a, x \rangle}, \quad (50)$$

$$F(x, y) = \frac{\phi\left((1 + \langle a, x \rangle)y - \langle a, y \rangle x\right)}{\left(1 + \langle a, x \rangle\right)^2}. \quad (51)$$

By Theorem 1.3, we know that $F(x, y)$ is a projective Finsler metric on its domain with $\mathbf{K} = 0$ and its projective factor is $P(x, y)$.

For the Finsler metric $F(x, y)$ in (51), the geodesic coefficients $G^i(x, y) = P(x, y)y^i$ are quadratic in $y \in \mathbb{R}^n$. Thus F is a Berwald metric. It is well-known that any Berwald metric with $\mathbf{K} = 0$ is locally Minkowskian. We conclude that the Finsler metric in (51) is locally Minkowskian.

Take a Randers norm, $\phi(y) = |y| + \langle b, y \rangle$, where $b \in \mathbb{R}^n$ is a vector with $|b| < 1$, we obtain a projective Randers metric with $\mathbf{K} = 0$,

$$F(x, y) = \frac{\sqrt{(1 + \langle a, x \rangle)^2 |y|^2 - 2(1 + \langle a, x \rangle)\langle a, y \rangle \langle x, y \rangle + \langle a, y \rangle^2 |x|^2}}{(1 + \langle a, x \rangle)^2} + \frac{(1 + \langle a, x \rangle)\langle b, y \rangle - \langle a, y \rangle \langle b, x \rangle}{(1 + \langle a, x \rangle)^2}.$$

This metric must be locally Minkowskian. This fact also follows from the main theorem in [Sh4].

7 $\mathbf{K} = 1$

In this section, we are going to construct projective Finsler metrics $F(x, y)$ with constant flag curvature $\mathbf{K} = 1$. Let $\psi(y)$ be a Minkowski norm on \mathbb{R}^n and $\varphi(y)$ a positively homogeneous function of degree on \mathbb{R}^n . We would like to find a projective Finsler metric $F(x, y)$ of constant flag curvature $\mathbf{K} = 1$ satisfying that $F(0, y) = \psi(y)$ and its projective factor $P(0, y) = \varphi(y)$. Suppose that there is a complex-valued solution to the following system

$$H_{x^k} = HH_{y^k} \quad (52)$$

with $H(0, y) = \varphi(y) + i\psi(y)$. We can express $H(x, y) = P(x, y) + iF(x, y)$. Plugging it into (52) yields

$$P_{x^k} - PP_{y^k} + FF_{y^k} + i\left\{F_{x^k} - (PF)_{y^k}\right\} = 0.$$

We obtain

$$F_{x^k} = (PF)_{y^k}, \quad P_{x^k} = PP_{y^k} - FF_{y^k}.$$

By Lemma 3.2, $F(x, y)$ is projective with constant flag curvature $\mathbf{K} = 1$ with $F(0, y) = \psi(y)$ and its projective factor is $P(x, y)$ with $P(0, y) = \varphi(y)$.

Let

$$H(x, y) := \sum_{m=0}^{\infty} \frac{1}{(m+1)!} \frac{d^m}{dt^m} \left[\left(\varphi(y+tx) + i\psi(y+tx) \right)^{m+1} \right]_{t=0}. \quad (53)$$

We assume that the above power series is convergent for $x \in \mathbb{R}^n$ close to the origin and any $y \in \mathbb{R}^n$. By a direct argument, one can verify that the function $H = H(x, y)$ in (53) satisfies equation (52) with $H(0, y) = \varphi(y) + i\psi(y)$.

Assume that $\phi(y) := \varphi(y) + i\psi(y)$ is y -analytic on $\mathbb{R}^n \setminus \{0\}$ so that for any $y \in \mathbb{R}^n \setminus \{0\}$, it can be extended to a function $\phi(y+zx)$, $z \in \mathbb{C}$ and $x \in \mathbb{R}^n$ such that there is a solution $H(x, y)$ to the following equation

$$H(x, y) = \phi\left(y + H(x, y)x\right). \quad (54)$$

Note that $H(0, y) = \phi(y)$. By the same argument as in Lemma 5.1, one can show that $H = H(x, y)$ satisfies equation (52) with $H(0, y) = \varphi(y) + i\psi(y)$.

Example 7.1 Let $\psi(y) = \cos(\alpha)|y|$ and $\varphi(y) = \sin(\alpha)|y|$ where $|\cdot|$ denotes the Euclidean norm on \mathbb{R}^n and α is an angle with $|\alpha| < \pi/2$. We have

$$\varphi(y) + i\psi(y) = ie^{-i\alpha}|y|.$$

Equation (54) becomes

$$H = ie^{-i\alpha} \sqrt{|y|^2 + 2\langle x, y \rangle H + |x|^2 H^2}. \quad (55)$$

We obtain

$$H = \frac{-\langle x, y \rangle + i\sqrt{(e^{2i\alpha} + |x|^2)|y|^2 - \langle x, y \rangle^2}}{e^{2i\alpha} + |x|^2}.$$

Express

$$H(x, y) = P(x, y) + iF(x, y).$$

Then $F(x, y)$ is a projective Finsler metric of constant flag curvature $\mathbf{K} = 1$ and $P(x, y)$ is the projective factor of $F(x, y)$. Let

$$\begin{aligned} A &:= \left(\cos(2\alpha)|y|^2 + (|x|^2|y|^2 - \langle x, y \rangle^2) \right)^2 + \left(\sin(2\alpha)|y|^2 \right)^2 \\ B &:= \cos(2\alpha)|y|^2 + (|x|^2|y|^2 - \langle x, y \rangle^2) \\ C &:= \sin(2\alpha)\langle x, y \rangle \\ C' &:= \left(\cos(2\alpha) + |x|^2 \right) \langle x, y \rangle, \\ D &:= |x|^4 + 2\cos(2\alpha)|x|^2 + 1. \end{aligned}$$

For an angle α with $0 \leq \alpha < \pi/2$,

$$\sqrt{(e^{2\alpha i} + |x|^2)|y|^2 - \langle x, y \rangle^2} = \sqrt{\frac{\sqrt{A+B}}{2}} + i\sqrt{\frac{\sqrt{A-B}}{2}}.$$

By an elementary argument we obtain

$$F(x, y) = \sqrt{\frac{\sqrt{A+B}}{2D} + \left(\frac{C}{D}\right)^2} + \frac{C}{D}, \quad (56)$$

$$P(x, y) = -\sqrt{\frac{\sqrt{A-B}}{2D} - \left(\frac{C}{D}\right)^2} - \frac{C'}{D}. \quad (57)$$

In dimension two, one can verify that the Finsler metric F in (56) is the Bryant metric [Br1][Br2].

We can generalize the above construction. Let $\phi(y)$ be an arbitrary Minkowski norm on \mathbb{R}^n and $\Theta(x, y)$ denote the Funk metric of ϕ . Let

$$\psi(y) := \cos(\alpha)\phi(y), \quad \varphi(y) := \sin(\alpha)\phi(y),$$

where α is an angle with $|\alpha| < \pi/2$. We have

$$\varphi(y) + i\psi(y) = ie^{-i\alpha}\phi(y).$$

Equation (54) becomes

$$H = ie^{-i\alpha}\phi(y + Hx). \quad (58)$$

Assume that the Funk metric $\Theta(x, y)$ is x -analytic in $x \in \mathbb{R}^n$ at $x = 0$ so that it can be extended to be a function $\Theta(zx, y)$ where $z \in \mathbb{C}$ with $|z| = 1$, $y \in \mathbb{R}^n$ and $x \in \mathbb{R}^n$ close to the origin. Then

$$H(x, y) := ie^{-i\alpha}\Theta\left(ie^{-i\alpha}x, y\right)$$

is a solution to (58) with $H(0, y) = ie^{-i\alpha}\phi(y) = \varphi(y) + i\psi(y)$. By the above argument, we know that the following function

$$F(x, y) := \frac{1}{2i}\left\{ie^{-i\alpha}\Theta\left(ie^{-i\alpha}x, y\right) - \overline{ie^{-i\alpha}\Theta\left(ie^{-i\alpha}x, y\right)}\right\} \quad (59)$$

is a projective Finsler metric with $\mathbf{K} = 1$ and its projector factor is given by

$$P(x, y) = \frac{1}{2}\left\{ie^{-i\alpha}\Theta\left(ie^{-i\alpha}x, y\right) + \overline{ie^{-i\alpha}\Theta\left(ie^{-i\alpha}x, y\right)}\right\}. \quad (60)$$

Further, $F(0, y) = \cos(\alpha)\phi(y)$ and $P(0, y) = \sin(\alpha)\phi(y)$.

References

- [AIM] P.L. Antonelli, R.S. Ingarden, and M. Matsumoto, *The theory of sprays and Finsler spaces with applications in physics and biology*, FTPH 58, Kluwer Academic Publishers, 1993.
- [Al] R. Alexander, *Planes for which the lines are the shortest paths between points*, Illinois J. of Math., **22**(1978), 177-190.
- [Alv1] J.C. Álvarez, *Contact topology, taut immersions, and Hilbert's fourth problem*, in "Differential and Symplectic Topology of Knots and Curves". S. Tabachnikov (Ed.). Adv. in Math. Sciences. AMS, 1999.
- [Alv2] J.C. Álvarez, *Symplectic geometry and Hilbert's fourth problem*, preprint.
- [AlFe] J.C. Álvarez and E. Fernandes, *Crofton formulas in projective Finsler spaces*, Electronic Research Announcements of the AMS. **4**(1998), 91-100.
- [AlGeSm] J.C. Álvarez, I.M. Gelfand and M. Smirnov, *Crofton Densities, symplectic geometry and Hilbert's fourth problem*, Arnold-Gelfand Mathematical Seminars, Geometry and Singularity Theory, V.I. Arnold, I.M. Gelfand, M. Smirnov, and V.S. Retakh (eds.). Birkhauser, Boston, 1997, pp. 77-92.
- [Am] R.V. Ambartzumian, *A note on pseudo-metrics on the plane*, Z. Wahrscheinlichkeitstheorie verw. Gebiete, **37**(1976), 145-155.
- [AZ] H. Akbar-Zadeh, *Sur les espaces de Finsler á courbures sectionnelles constantes*, Bull. Acad. Roy. Bel. Cl, Sci, 5e Série - Tome LXXXIV (1988), 281-322.
- [BaRo] D. Bao and C. Robles, *On Randers metrics of constant curvature*, Reports on Mathematical Physics (to appear).
- [BaSh] D. Bao and Z. Shen, *Finsler metrics of constant curvature on the Lie group S^3* , Journal of London Mathematical Society (to appear).
- [BCS] D. Bao, S.S. Chern and Z. Shen, *An introduction to Riemann-Finsler geometry*, Springer, 2000.
- [Be1] L. Berwald, *Über eine characteristic Eigenschaft der allgemeinen Räume konstanter Krümmung mit gradlinigen Extremalen*, Monatsh. Math. Phys. **36**(1929), 315-330.
- [Be2] L. Berwald, *Über die n-dimensionalen Geometrien konstanter Krümmung, in denen die Geraden die kürzesten sind*, Math. Z. **30**(1929), 449-469.

- [Bl] W. Blaschke, *Integral geometrie 11: Zur Variationsrechnung*, Abh. Math. Sem. Univ. Hamburg, **11**(1936), 359-366.
- [Br1] R. Bryant, *Finsler structures on the 2-sphere satisfying $K = 1$* , Finsler Geometry, Contemporary Mathematics **196**, Amer. Math. Soc., Providence, RI, 1996, 27-42.
- [Br2] R. Bryant, *Projectively flat Finsler 2-spheres of constant curvature*, Selecta Math., New Series, **3**(1997), 161-204.
- [Br3] R. Bryant, *Some remarks on Finsler manifolds with constant flag curvature*, Houston J. of Math. **28**(2002), 221-262.
- [Bu] H. Busemann, *Problem IV: Desarguesian spaces*, in Mathematical Developments arising from Hilbert Problems, Proc. Symp. Pure Math. **28**(1976), Amer. Math. Soc., Providence, RI, 131-141.
- [Fk1] P. Funk, *Über Geometrien, bei denen die Geraden die Kürzesten sind*, Math. Annalen **101**(1929), 226-237.
- [Fk2] P. Funk, *Über zweidimensionale Finslersche Räume, insbesondere über solche mit geradlinigen Extremalen und positiver konstanter Krümmung*, Math. Zeitschr. **40**(1936), 86-93.
- [Fk3] P. Funk, *Eine Kennzeichnung der zweidimensionalen elliptischen Geometrie*, Österreichische Akad. der Wiss. Math., Sitzungsberichte Abteilung II **172**(1963), 251-269.
- [Ha] G. Hamel, *Über die Geometrien, in denen die Geraden die Kürzesten sind*, Math. Ann. **57**(1903), 231-264.
- [Hi] D. Hilbert, *Mathematical Problems*, Bull. of Amer. Math. Soc. **37**(2001), 407-436. Reprinted from Bull. Amer. Math. Soc. **8** (July 1902), 437-479.
- [MaSh] M. Matsumoto and H. Shimada, *The corrected fundamental theorem on the Randers spaces of constant curvature*, Tensor, N.S. (to appear).
- [Ok] T. Okada, *On models of projectively flat Finsler spaces of constant negative curvature*, Tensor, N. S. **40**(1983), 117-123.
- [Po] A.V. Pogorelov, *Hilbert's Fourth Problem*, Winston & Wiley, New York, 1982.
- [Ran] G. Randers, *On an asymmetric metric in the four-space of general relativity*, Phys. Rev. **59**(1941), 195-199.
- [Rap] A. Rapcsák, *Über die bahntreuen Abbildungen metrischer Räume*, Publ. Math. Debrecen, **8**(1961), 285-290.

- [Sh1] Z. Shen, *Differential Geometry of Spray and Finsler Spaces*, Kluwer Academic Publishers, Dordrecht, 2001.
- [Sh2] Z. Shen, *Finsler metrics with $\mathbf{K} = 0$ and $\mathbf{S} = 0$* , Canadian J. Math. (to appear).
- [Sh3] Z. Shen, *Two-dimensional Finsler metrics of constant flag curvature*, Manuscripta Mathematica (to appear).
- [Sh4] Z. Shen, *Projectively flat Randers metrics of constant flag curvature*, Math. Ann. (to appear).
- [Sz] Z.I. Szabó, *Hilbert's fourth problem, I*, Adv. in Math. **59**(1986), 185-301.

Math Dept, IUPUI, 402 N. Blackford Street, Indianapolis, IN 46202-3216, USA.
zshen@math.iupui.edu