

Correlations Between Zeros of a Random Polynomial

Pavel Bleher¹ and Xiaojun Di¹

Received November 13, 1996

We obtain exact analytical expressions for correlations between real zeros of the Kac random polynomial. We show that the zeros in the interval $(-1, 1)$ are asymptotically independent of the zeros outside of this interval, and that the straightened zeros have the same limit-translation-invariant correlations. Then we calculate the correlations between the straightened zeros of the $O(1)$ random polynomial.

KEY WORDS: Real random polynomials; correlations between zeros; scaling limit; determinants of block matrices.

1. INTRODUCTION

Let $f_n(t)$ be a real random polynomial of degree n ,

$$f_n(t) = c_0 + c_1 t + \dots + c_n t^n \tag{1.1}$$

where c_0, c_1, \dots, c_n are independent real random variables. Distribution of zeros for various classes of random polynomials is studied in the classical papers by Bloch and Polya [BP], Littlewood and Offord [LO], Erdős and Offord [EO], Erdős and Turán [ET], and Kac [K1-K3]. We will assume that the coefficients c_0, c_1, \dots, c_n are normally distributed with

$$E c_j = 0, \quad E c_j^2 = \sigma_j^2 \tag{1.2}$$

In the case when

$$\sigma_j^2 = 1$$

¹ Department of Mathematical Sciences, Indiana University-Purdue University at Indianapolis, Indianapolis, Indiana 46202; e-mail: bleher@math.iupui.edu, xdi@math.iupui.edu.

$f_n(t)$ is the Kac random polynomial. Another interesting case is when

$$\sigma_j^2 = \binom{n}{j}$$

As is pointed out by Edelman and Kostlan [EK], “this particular random polynomial is probably the more natural definition of a random polynomial.” We call this polynomial the $O(1)$ random polynomial because its m -point joint probability distribution of zeros is $O(1)$ -invariant for all m (see Section 5 below). The $O(1)$ random polynomial can be viewed as the Majorana spin state [Maj] with real random coefficients, and it models a chaotic spin wavefunction in the Majorana representation. See the papers by Leboeuf [Leb1, Leb2], Leboeuf and Shukla [LS], Bogomolny, Bohigas, and Leboeuf [BBL2], and Hannay [Han], where the $SU(2)$ and some other random polynomials are introduced and studied, that represent the Majorana spin states with complex random coefficients.

Let $\{\tau_1, \dots, \tau_k\}$ be the set of real zeros of $f_n(t)$. Consider the distribution function of the real zeros,

$$P_n(t) = \mathbb{E} \# \{j: \tau_j \leq t\}$$

where the mathematical expectation is taken with respect to the joint distribution of the coefficients c_0, \dots, c_n . Let

$$p_n(t) = P'_n(t)$$

be the density function. By the Kac formula (see, e.g., [K3]),

$$p_n(t) = \frac{\sqrt{A_n(t) C_n(t) - B_n^2(t)}}{\pi A_n(t)} \quad (1.3)$$

where

$$\begin{aligned} A_n(t) &= \sum_{j=0}^n \sigma_j^2 t^{2j} \\ B_n(t) &= \sum_{j=1}^n j \sigma_j^2 t^{2j-1} = \frac{A'_n(t)}{2} \\ C_n(t) &= \sum_{j=1}^n j^2 \sigma_j^2 t^{2j-2} = \frac{A''_n(t)}{4} + \frac{A'_n(t)}{4t} \end{aligned} \quad (1.4)$$

The derivation of (1.3) by Kac is rather complex. A short proof of (1.3) is given in the paper [EK] by Edelman and Kostlan. See also the papers by

Hannay [Han] and Mesincescu, Bessis, Fournier, Mantica, and Aaron [M-A], and Section 2 below. The formula (1.3) implies that for the Kac random polynomial,

$$\lim_{n \rightarrow \infty} p_n(t) = p(t) = \frac{1}{\pi |1 - t^2|}, \quad t \neq \pm 1 \tag{1.5}$$

and

$$p_n(\pm 1) = \frac{1}{\pi} \left[\frac{n(n+2)}{12} \right]^{1/2}$$

(see [K3], [BS], and [EK]). The limiting density $p(t)$ is not integrable at ± 1 , and this means that the zeros are mostly located near ± 1 . Observe, in addition, that $p_n(t)$ is an even function of t , and the distribution $p_n(t) dt$ is invariant with respect to the transformation $t \rightarrow 1/t$. Kac [K1] proves that the expected number of real zeros has the asymptotics

$$N_n = \int_{-\infty}^{\infty} p_n(t) dt = (2/\pi) \log n + O(1)$$

Kac [K2], Erdős and Offord [EO], Stevens [Ste], Ibragimov and Maslova [IM], Logan and Shepp [LS], Edelman and Kostlan [EK], and others extend this asymptotics to various classes of the random coefficients $\{c_j\}$. Maslova [Mas1] evaluates the variance of the number of real zeros as

$$\text{Var } \#\{j: f_n(\tau_j) = 0\} = \frac{4}{\pi} \left(1 - \frac{2}{\pi} \right) \ln n(1 + o(1)), \quad n \rightarrow \infty$$

and she proves the central limit theorem for the number of real zeros (see [Mas2]), for a class of distributions of the random coefficients $\{c_j\}$.

In this paper we are interested in correlations between the zeros τ_j of the Kac random polynomial. Let us consider first the zeros in the interval $(-1, 1)$. Define straightening of τ_j as

$$\zeta_j = P(\tau_j), \quad P(t) = \int_0^t p(u) du$$

In the limit when $n \rightarrow \infty$, the straightened zeros ζ_j are uniformly distributed on the real line, so that

$$\lim_{n \rightarrow \infty} E \#\{j: a < \zeta_j \leq b\} = b - a \tag{1.6}$$

From (1.5) we get that

$$P(t) = \int_0^t \frac{du}{\pi(1-u^2)} = \frac{1}{2\pi} \ln \left| \frac{1+t}{1-t} \right| = \frac{1}{\pi} \operatorname{artanh} t$$

hence

$$\zeta_j = \frac{1}{\pi} \operatorname{artanh} \tau_j \tag{1.7}$$

Let $p_{nm}(s_1, \dots, s_m)$ be the joint probability distribution density of the straightened zeros ζ_j ,

$$p_{nm}(s_1, \dots, s_m) = \lim_{\Delta s_1, \dots, \Delta s_m \rightarrow 0} \frac{\Pr\{\exists \zeta_{j_1} \in [s_1, s_1 + \Delta s_1], \dots, \exists \zeta_{j_m} \in [s_m, s_m + \Delta s_m]\}}{|\Delta s_1 \cdots \Delta s_m|} \tag{1.8}$$

It coincides with the correlation function

$$k_{nm}(s_1, \dots, s_m) = \lim_{\Delta s_1, \dots, \Delta s_m \rightarrow 0} \frac{E[\xi_n(s_1, s_1 + \Delta s_1) \cdots \xi_n(s_m, s_m + \Delta s_m)]}{|\Delta s_1 \cdots \Delta s_m|} \tag{1.9}$$

where

$$\xi_n(a, b) = \#\{j: a < \zeta_j \leq b\}$$

Our aim is to find the limit correlation functions

$$k_m(s_1, \dots, s_m) = \lim_{n \rightarrow \infty} k_{nm}(s_1, \dots, s_m) \tag{1.10}$$

We prove the following results.

Theorem 1.1. The limit two-point correlation function $k_2(s_1, s_2)$ of the straightened zeros $\zeta_j = \pi^{-1} \operatorname{artanh} \tau_j$ of the Kac random polynomial is equal to

$$k_2(s_1, s_2) = \tanh^2 \pi(s_1 - s_2) + \frac{|\sinh \pi(s_1 - s_2)|}{\cosh^2 \pi(s_1 - s_2)} \arcsin \frac{1}{\cosh \pi(s_1 - s_2)} \tag{1.11}$$

Observe that $k_2(s_1, s_2)$ depends only on $s_1 - s_2$, and it has the following asymptotics:

$$k_2(s_1, s_2) = \frac{\pi^2}{2} |s_1 - s_2| + O(|s_1 - s_2|^2), \quad |s_1 - s_2| \rightarrow 0$$

$$k_2(s_1, s_2) = 1 - \frac{16}{3} e^{-4\pi |s_1 - s_2|} + O(e^{-6\pi |s_1 - s_2|}), \quad |s_1 - s_2| \rightarrow \infty$$

The graph of $k_2(0, s)$ is given in Fig. 1.

Theorem 1.2. The limit m -point correlation function $k_m(s_1, \dots, s_m)$ of the straightened zeros $\zeta_j = \pi^{-1} \operatorname{artanh} \tau_j$ is equal to

$$k_m(s_1, \dots, s_m) = 2^{-m} \prod_{1 \leq i < j \leq m} \tanh^4 \pi(s_i - s_j) \times \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| e^{-1/2(Y\Gamma_m, Y)} dy_1 \dots dy_m \quad (1.12)$$

where $Y = (y_1, \dots, y_m)$ and the matrix Γ_m is defined as

$$\Gamma_m = \left(\frac{1}{\cosh \pi(s_i - s_j)} \right)_{i, j=1}^m \quad (1.13)$$

In particular, $k_m(s_1, \dots, s_m)$ depends only on the differences of s_1, \dots, s_m , hence it is translation invariant.

The proof of Theorems 1.1 and 1.2 is given in Sections 2, 3 and 4 below. It is based on computation of the determinant of some matrices




Fig. 1. The two-point correlation function of straightened zeros of the Kac random polynomial.

which consist of 2×2 blocks. This computation is of independent interest. The basic example is the matrix

$$\Delta_m = \begin{pmatrix} \Delta_{11} & \Delta_{12} & \cdots & \Delta_{1m} \\ \Delta_{21} & \Delta_{22} & \cdots & \Delta_{2m} \\ \cdots & \cdots & \cdots & \cdots \\ \Delta_{m1} & \Delta_{m2} & \cdots & \Delta_{mm} \end{pmatrix} \tag{1.14}$$

where

$$\Delta_{ij} = \begin{pmatrix} \frac{1}{1-t_it_j} & \frac{t_i}{(1-t_it_j)^2} \\ \frac{t_j}{(1-t_it_j)^2} & \frac{1+t_it_j}{(1-t_it_j)^3} \end{pmatrix}, \quad i, j = 1, \dots, m \tag{1.15}$$

We prove in Section 4 that

$$\det \Delta_m = \frac{\prod_{1 \leq i < j \leq m} (t_i - t_j)^8}{\prod_{i=1}^m (1-t_i^2)^4 \prod_{1 \leq i < j \leq m} (1-t_it_j)^8} \tag{1.16}$$

It is interesting to note that determinants of matrices consisting of 2×2 blocks appear also in the theory of random matrices (see, e.g., [Dys] and [Meh]), statistical physics, and other applications.

Consider now zeros τ_j with $|\tau_j| > 1$. Define straightening of τ_j as

$$\zeta_j = P(\tau_j), \quad P(t) = \begin{cases} \int_t^{-\infty} p(u) du & \text{if } t < -1 \\ \int_t^{\infty} p(u) du & \text{if } t > 1 \end{cases} \tag{1.17}$$

In the limit when $n \rightarrow \infty$, the straightened zeros ζ_j are uniformly distributed on the real line. From (1.5)

$$P(t) = \frac{1}{2\pi} \ln \left| \frac{1+t}{1-t} \right| = \frac{1}{\pi} \operatorname{artanh} t^{-1} \tag{1.18}$$

so that

$$\zeta_j = \pi^{-1} \operatorname{artanh} \tau_j^{-1} \tag{1.19}$$

Denote by $k_{nm}^{\text{out}}(s_1, \dots, s_m)$ the correlation function of the straightened zeros ζ_j with $|\tau_j| > 1$.

Theorem 1.3.

$$k_{nm}^{\text{out}}(s_1, \dots, s_m) = k_{nm}(s_1, \dots, s_m) \tag{1.20}$$

In other words, the correlation functions of the straightened zeros outside of the interval $(-1, 1)$ coincide with those inside of the interval. Finally, let us consider correlation between zeros inside of the interval $(-1, 1)$ and outside of this interval. Let $K_{nm}(t_1, \dots, t_m)$ be the correlation function of the zeros τ_j (without straightening).

Theorem 1.4. Assume that $|t_1|, \dots, |t_l| < 1$ and $|t_{l+1}|, \dots, |t_m| > 1$. Then the limit

$$\lim_{n \rightarrow \infty} K_{nm}(t_1, \dots, t_m) = K_m(t_1, \dots, t_m) \tag{1.21}$$

exists and

$$K_m(t_1, \dots, t_m) = K_l(t_1, \dots, t_l) K_{m-l}(t_{l+1}, \dots, t_m) \tag{1.22}$$

This means that the zeros inside and outside of the interval $(-1, 1)$ are asymptotically independent. Observe that

$$k_m(s_1, \dots, s_m) = \left[\frac{K_m(t_1, \dots, t_m)}{p(t_1) \cdots p(t_m)} \right]_{t_1 = P^{-1}(s_1), \dots, t_m = P^{-1}(s_m)} \tag{1.23}$$

provided that either all $|t_j| < 1$ or all $|t_j| > 1$ (cf. the formula (2.14) below). Proof of Theorems 1.3 and 1.4 is given in the end of Section 4.

In Sections 5 and 6 we investigate correlation functions of real zeros of the $O(1)$ random polynomial.

2. GENERAL FORMULAE

Let

$$f_n(t) = \sum_{j=0}^n c_j t^j \tag{2.1}$$

be a polynomial whose coefficients c_j are random variables with an absolutely continuous joint distribution. Let

$$\xi_n(a, b) = \# \{ \tau_k : a < \tau_k \leq b, f_n(\tau_k) = 0 \} \tag{2.2}$$

be the number of real roots of $f_n(t)$ between a and b , and let $p_n(t)$ be the density of real zeros t_k of $f_n(t)$, so that

$$E \xi_n(a, b) = \int_a^b p_n(t) dt \tag{2.3}$$

It is not difficult to show that

$$p_n(t) = \int_{-\infty}^{\infty} |y| D_n(0, y; t) dy \tag{2.4}$$

where $D_n(x, y; t)$ is a joint distribution density of $f_n(t)$ and $f'_n(t)$,

$$\Pr\{a < f_n(t) \leq b; c < f'_n(t) \leq d\} = \int_a^b \int_c^d D_n(x, y; t) dx dy \tag{2.5}$$

Indeed, if $f'_n(t) = y$ then asymptotically as $\Delta t \rightarrow 0$, the function $f_n(t)$ has a zero in the interval $[t, t + \Delta t]$ if $f_n(t)$ is in the interval $[0, -y \Delta t]$, and this gives (2.5). Similarly, the m -point correlation function $K_{nm}(t_1, \dots, t_m)$ for pairwise different t_1, \dots, t_m is equal to

$$\begin{aligned} &K_{nm}(t_1, \dots, t_m) \\ &= \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| D_{nm}(0, y_1, \dots, 0, y_m; t_1, \dots, t_m) dy_1 \dots dy_m \end{aligned} \tag{2.6}$$

where $D_{nm}(x_1, y_1, \dots, x_m, y_m; t_1, \dots, t_m)$ is a joint distribution density of the vector

$$F_n = (f_n(t_1), f'_n(t_1), \dots, f_n(t_m), f'_n(t_m))$$

so that

$$\begin{aligned} &\Pr\{a_1 < f_n(t_1) \leq b_1; c_1 < f'_n(t_1) \leq d_1; \dots; a_m < f_n(t_m) \leq b_m; c_m < f'_n(t_m) \leq d_m\} \\ &= \int_{a_1}^{b_1} \int_{c_1}^{d_1} \dots \int_{a_m}^{b_m} \int_{c_m}^{d_m} D_{nm}(x_1, y_1, \dots, x_m, y_m; t_1, \dots, t_m) dx_1 dy_1 \dots dx_m dy_m \end{aligned} \tag{2.7}$$

If $\{c_j\}$ are independent random variables with $\text{Var } c_j > 0$ then the covariance matrix of the vector F_n is positive, provided that $n \geq 2m - 1$ (see Appendix B at the end of the paper). Similar formulae are derived for the correlation functions of complex zeros of random polynomials with complex and real coefficients (see [Han] and [M-A]).

Observe that

$$E \prod_{j=1}^m \xi_n(a_j, b_j) = \int_{a_1}^{b_1} \dots \int_{a_m}^{b_m} K_{nm}(t_1, \dots, t_m) dt_1 \dots dt_m \tag{2.8}$$

provided that $(a_1, b_1), \dots, (a_m, b_m)$ are pairwise disjoint, and

$$p_n(t) = K_{n1}(t), \quad E(\xi_n(a, b)) = \int_a^b K_{n1}(t) dt \quad (2.9)$$

For the general case, when $(a_1, b_1), \dots, (a_m, b_m)$ may intersect, we have the following extension of (2.8):

$$E \prod_{j=1}^m \xi_n(a_j, b_j) = \sum_{(A_1, \dots, A_l)} \prod_{j=1}^l \left(\int_{\bigcap_{i \in A_j} (a_i, b_i)} dt_j \right) K_{nl}(t_1, \dots, t_l) \quad (2.10)$$

where the sum is taken over all possible partitions (A_1, \dots, A_l) of $\{1, \dots, m\}$, such that

$$\begin{aligned} A_i \cap A_j &= \emptyset, \quad i \neq j \\ A_1 \cup \dots \cup A_l &= \{1, \dots, m\} \\ |A_i| &\geq 1, \quad i = 1, \dots, l \end{aligned} \quad (2.11)$$

In particular, when $m = 2$ we have

$$E[\xi_n(a_1, b_1) \xi_n(a_2, b_2)] = \int_{a_1}^{b_1} \int_{a_2}^{b_2} K_{n2}(t_1, t_2) dt_1 dt_2 \quad (2.12)$$

if $(a_1, b_1) \cap (a_2, b_2) = \emptyset$, and

$$E[\xi_n^2(a, b)] = \int_a^b p_n(t) dt + \int_a^b \int_a^b K_{n2}(t_1, t_2) dt_1 dt_2 \quad (2.13)$$

From the definition (1.9) of the m -point correlation function, it follows that the m -point correlation function $k_{nm}(s_1, \dots, s_m)$ of the straightened zeros $\zeta_j = P(\tau_j)$ is related to the m -point correlation function $K_{nm}(t_1, \dots, t_m)$ of the zeros τ_j by the formula

$$k_{nm}(s_1, \dots, s_m) = \left[\frac{K_{nm}(t_1, \dots, t_m)}{p(t_1) \cdots p(t_m)} \right]_{t_1 = P^{-1}(s_1), \dots, t_m = P^{-1}(s_m)} \quad (2.14)$$

Assume now that the coefficients c_j are independent Gaussian variables with zero mean and the variances σ_j^2 , $j = 0, \dots, n$. Then $D_{n1}(x, y; t)$ is a Gaussian distribution density with the covariance matrix

$$A = \begin{pmatrix} E f_n^2(t) & E f_n(t) f_n'(t) \\ E f_n(t) f_n'(t) & E (f_n'(t))^2 \end{pmatrix} = \begin{pmatrix} A_n(t) & B_n(t) \\ B_n(t) & C_n(t) \end{pmatrix} \quad (2.15)$$

where $A_n(t)$, $B_n(t)$ and $C_n(t)$ are defined in (1.4), and from (2.4) we get the Kac formula (1.3).

3. TWO-POINT CORRELATION FUNCTION FOR THE KAC POLYNOMIAL

Let $f_n(t) = c_0 + c_1 t + \dots + c_n t^n$ be the Kac polynomial, so that c_k , $k = 0, \dots, n$, are real independent Gaussian random variables with

$$E c_k = 0, \quad E c_k^2 = 1 \tag{3.1}$$

Consider the covariance matrix A_n of the Gaussian vector $(f_n(t_1), f'_n(t_1), f_n(t_2), f'_n(t_2))$. From (3.1)

$$\begin{aligned} E f_n(t_1) f_n(t_2) &= \sum_{k=0}^n (t_1 t_2)^k = \frac{1 - (t_1 t_2)^{n+1}}{1 - t_1 t_2} \\ E f'_n(t_1) f_n(t_2) &= \frac{\partial}{\partial t_1} \left[\frac{1 - (t_1 t_2)^{n+1}}{1 - t_1 t_2} \right] \\ E f'_n(t_1) f'_n(t_2) &= \frac{\partial^2}{\partial t_1 \partial t_2} \left[\frac{1 - (t_1 t_2)^{n+1}}{1 - t_1 t_2} \right] \end{aligned} \tag{3.2}$$

Assume that $|t_1|, |t_2| < 1$. Then from (3.2) we obtain that

$$\lim_{n \rightarrow \infty} A_n = A \tag{3.3}$$

with

$$A = \begin{pmatrix} \frac{1}{1-t_1^2} & \frac{t_1}{(1-t_1^2)^2} & \frac{1}{1-t_1 t_2} & \frac{t_1}{(1-t_1 t_2)^2} \\ \frac{t_1}{(1-t_1^2)^2} & \frac{1+t_1^2}{(1-t_1^2)^3} & \frac{t_2}{(1-t_1 t_2)^2} & \frac{1+t_1 t_2}{(1-t_1 t_2)^3} \\ \frac{1}{1-t_1 t_2} & \frac{1+t_1 t_2}{(1-t_1 t_2)^3} & \frac{1}{1-t_2^2} & \frac{t_2}{(1-t_2^2)^2} \\ \frac{t_1}{(1-t_1 t_2)^2} & \frac{1+t_1 t_2}{(1-t_1 t_2)^3} & \frac{t_2}{(1-t_2^2)^2} & \frac{1+t_2^2}{(1-t_2^2)^3} \end{pmatrix} \tag{3.4}$$

We prove in the Section 4 below that

$$\det A = \frac{(t_1 - t_2)^8}{(1 - t_1^2)^4 (1 - t_2^2)^4 (1 - t_1 t_2)^8} \tag{3.5}$$

Let Ω be the two-by-two matrix obtained by removing the first and the third rows and columns from A^{-1} . Then

$$\Omega = \begin{pmatrix} A & B \\ B & C \end{pmatrix} \tag{3.6}$$

where

$$\begin{aligned} A &= (1 - t_1 t_2)^4 (1 - t_1^2)^3 / (t_1 - t_2)^4 \\ B &= (1 - t_1 t_2)^3 (1 - t_1^2)^2 (1 - t_2^2)^2 / (t_1 - t_2)^4 \\ C &= (1 - t_1 t_2)^4 (1 - t_2^2)^3 / (t_1 - t_2)^4 \end{aligned} \tag{3.7}$$

By (2.6), the correlation function $K_2(t_1, t_2)$ is equal to

$$K_2(t_1, t_2) = \frac{1}{4\pi^2 \sqrt{\det A}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(Y\Omega, Y)} dy_1 dy_2 \tag{3.8}$$

where $Y = (y_1, y_2)$. Since

$$\begin{aligned} &\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(Ay_1^2 + 2By_1 y_2 + Cy_2^2)} dy_1 dy_2 \\ &= \frac{4}{AC(1 - \delta^2)} \left(1 + \frac{\delta}{\sqrt{1 - \delta^2}} \arcsin \delta \right), \quad \delta = \frac{B}{\sqrt{AC}} \end{aligned} \tag{3.9}$$

(see Appendix A), we obtain that

$$\begin{aligned} K_2(t_1, t_2) &= \frac{(t_1 - t_2)^2}{\pi^2(1 - t_1 t_2)^2 (1 - t_1^2)(1 - t_2^2)} \\ &\quad + \frac{|t_1 - t_2|}{\pi^2(1 - t_1 t_2)^2 \sqrt{(1 - t_1^2)(1 - t_2^2)}} \\ &\quad \times \arcsin \frac{\sqrt{(1 - t_1^2)(1 - t_2^2)}}{1 - t_1 t_2} \end{aligned} \tag{3.10}$$

Consider the correlation function $k_2(s_1, s_2)$ of the straightened zeroes $\zeta_j = \pi^{-1} \operatorname{artanh} \tau_j$. By (2.14),

$$k_2(s_1, s_2) = \frac{K_2(t_1, t_2)}{p(t_1)p(t_2)}, \quad t_1 = \tanh(\pi s_1), \quad t_2 = \tanh(\pi s_2) \tag{3.11}$$

Since

$$p(t) = \frac{1}{\pi(1-t^2)}$$

(see (1.5)), we obtain that

$$\begin{aligned} k_2(s_1, s_2) &= \frac{(t_1 - t_2)^2}{(1 - t_1 t_2)^2} + \frac{|t_1 - t_2| \sqrt{(1 - t_1^2)(1 - t_2^2)}}{(1 - t_1 t_2)^2} \arcsin \frac{\sqrt{(1 - t_1^2)(1 - t_2^2)}}{1 - t_1 t_2} \\ &= \tanh^2 \pi(s_1 - s_2) + \frac{|\sinh \pi(s_1 - s_2)|}{\cosh^2 \pi(s_1 - s_2)} \arcsin \frac{1}{\cosh \pi(s_1 - s_2)} \end{aligned} \quad (3.12)$$

Theorem 1.1 is proved.

4. HIGHER ORDER CORRELATION FUNCTIONS FOR THE KAC POLYNOMIAL

Let $f_n(t)$ be the Kac polynomial, and let t_1, t_2, \dots, t_m be $m \geq 3$ distinct points in the interval $(-1, 1)$. Denote by $\Delta_m^{(n)}$ the covariance matrix of the Gaussian vector

$$(f_n(t_1), f'_n(t_1), \dots, f_n(t_m), f'_n(t_m))$$

and by Δ_m the limit of $\Delta_m^{(n)}$ as $n \rightarrow \infty$,

$$\Delta_m = \lim_{n \rightarrow \infty} \Delta_m^{(n)} \quad (4.1)$$

Then

$$\Delta_m^{(n)} = \begin{pmatrix} \Delta_{11}^{(n)} & \Delta_{12}^{(n)} & \cdots & \Delta_{1m}^{(n)} \\ \Delta_{21}^{(n)} & \Delta_{22}^{(n)} & \cdots & \Delta_{2m}^{(n)} \\ \cdots & \cdots & \cdots & \cdots \\ \Delta_{m1}^{(n)} & \Delta_{m2}^{(n)} & \cdots & \Delta_{mm}^{(n)} \end{pmatrix} \quad (4.2)$$

where

$$\Delta_{ij}^{(n)} = \begin{pmatrix} \mathbf{E} f_n(t_i) f_n(t_j) & \mathbf{E} f_n(t_i) f'_n(t_j) \\ \mathbf{E} f'_n(t_i) f_n(t_j) & \mathbf{E} f'_n(t_i) f'_n(t_j) \end{pmatrix} \quad (4.3)$$

and by (3.2),

$$A_m = \begin{pmatrix} \Delta_{11} & \Delta_{12} & \cdots & \Delta_{1m} \\ \Delta_{21} & \Delta_{22} & \cdots & \Delta_{2m} \\ \cdots & \cdots & \cdots & \cdots \\ \Delta_{m1} & \Delta_{m2} & \cdots & \Delta_{mm} \end{pmatrix} \tag{4.4}$$

where

$$\Delta_{ij} = \begin{pmatrix} \frac{1}{1-t_i t_j} & \frac{t_i}{(1-t_i t_j)^2} \\ \frac{t_j}{(1-t_i t_j)^2} & \frac{1+t_i t_j}{(1-t_i t_j)^3} \end{pmatrix} \tag{4.5}$$

[cf. (3.4)].

If Ω_m denotes the $m \times m$ matrix obtained by removing all the odd number rows and columns from A_m^{-1} , then by (2.6), the correlation function $K_m(t_1, \dots, t_m)$ is equal to

$$K_m(t_1, \dots, t_m) = \frac{1}{(2\pi)^m \sqrt{\det A_m}} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} |y_1 \cdots y_m| e^{-1/2(Y\Omega_m Y)} dy_1 \cdots dy_m \tag{4.6}$$

where $Y = (y_1, \dots, y_m)$. We have the following extension of the formula (3.6).

Proposition 4.1.

$$\det A_m = \frac{\prod_{1 \leq i < j \leq m} (t_i - t_j)^8}{\prod_{i=1}^m (1 - t_i^2)^4 \prod_{1 \leq i < j \leq m} (1 - t_i t_j)^8} \tag{4.7}$$

The proof of Proposition 4.1 uses the following lemma.

Lemma 4.2. Let $f_n(t)$ ($n \geq 3$) be any random polynomial and t_1, \dots, t_m be any m real numbers. Let $A_m^{(n)}$ be the covariance matrix of the Gaussian random vector

$$(f_n(t_1), f'_n(t_1), \dots, f_n(t_m), f'_n(t_m))$$

which is defined in (4.2) and (4.3). Then

$$\det A_m^{(n)} = P_n(t_1, \dots, t_m) \prod_{1 \leq i < j \leq m} (t_i - t_j)^8 \tag{4.8}$$

where $P_n(t_1, \dots, t_m)$ is a polynomial.

Proof. To simplify notation we drop the indices m, n in the matrix $A_m^{(n)}$. We have

$$A = (A_{ij})_{i,j=1,\dots,m} \quad (4.9)$$

where

$$A_{ij} = \begin{pmatrix} \mathbb{E} f_n(t_i) f_n(t_j) & \mathbb{E} f_n(t_i) f'_n(t_j) \\ \mathbb{E} f'_n(t_i) f_n(t_j) & \mathbb{E} f'_n(t_i) f'_n(t_j) \end{pmatrix} \quad (4.10)$$

In the following discussion we consider linear transformations of the matrix A which do not change its determinant. By subtracting the first and second column of A_{i1} from the first and second column of A_{ij} , respectively, we get the matrix $A^{(1)}$ with the 2×2 blocks

$$A_{ij}^{(1)} = \begin{pmatrix} \mathbb{E} f_n(t_i)(f_n(t_j) - f_n(t_1)) & \mathbb{E} f_n(t_i)(f'_n(t_j) - f'_n(t_1)) \\ \mathbb{E} f'_n(t_i)(f_n(t_j) - f_n(t_1)) & \mathbb{E} f'_n(t_i)(f'_n(t_j) - f'_n(t_1)) \end{pmatrix} \quad (4.11)$$

Since $f_n(t)$ is a polynomial, we can take the factor $(t_j - t_1)$ out of the first two columns of the matrix $A^{(1)}$, and this proves that $\det A$ is divisible by $(t_j - t_1)^2$. Repeating the same operation on rows we get the factor $(t_j - t_1)^4$. How to get $(t_j - t_1)^8$? To that end let us subtract the second column of the matrix $A_{i1}^{(1)} = A_{i1}$ multiplied by $(t_j - t_1)$ from the first column of the matrix $A_{ij}^{(1)}$. This produces the matrix

$$A_{ij}^{(2)} = \begin{pmatrix} \mathbb{E} f_n(t_i)[f_n(t_j) - f_n(t_1) - (t_j - t_1)f'_n(t_1)] & \mathbb{E} f_n(t_i)[f'_n(t_j) - f'_n(t_1)] \\ \mathbb{E} f'_n(t_i)[f_n(t_j) - f_n(t_1) - (t_j - t_1)f'_n(t_1)] & \mathbb{E} f'_n(t_i)[f'_n(t_j) - f'_n(t_1)] \end{pmatrix} \quad (4.12)$$

Now we can take $(t_j - t_1)^2$ out of the first column and $(t_j - t_1)$ out of the second column of the matrix $A^{(2)}$. Repeating the same operations over the rows we get that $\det A$ is divisible by $(t_j - t_1)^6$. Finally, let us observe that by the Taylor formula

$$f_n(t_j) - f_n(t_1) - (t_j - t_1) f'_n(t_1) = \frac{(t_j - t_1)^2}{2} f''_n(t_1) + O(|t_j - t_1|^3)$$

and

$$f'_n(t_j) - f'_n(t_1) = (t_j - t_1) f''_n(t_1) + O(|t_j - t_1|^2)$$

hence if we subtract the second column of the matrix $A_{ij}^{(2)}$ multiplied by $(t_j - t_1)/2$ from its first column, the difference is of the order of $|t_j - t_1|^3$,

and we can take the factor $(t_j - t_1)^3$ out of the first column and $(t_j - t_1)$ out of the second column. This gives the factor $(t_j - t_1)^4$. The same factor is taken out of the rows, hence $\det \Delta$ is divisible by $(t_j - t_1)^8$. Similarly, it is divisible by $(t_j - t_i)^8$ for all $i \neq j$, and hence it is divisible by their product. Lemma 4.2 is proved.

Proof of Proposition 4.1. By (4.1), we have

$$\det \Delta_m = \lim_{n \rightarrow \infty} \det \Delta_m^{(n)} \tag{4.13}$$

for all t_1, \dots, t_m in the interval $(-1, 1)$. In fact, the limit (4.13) holds for all complex t_1, \dots, t_m in the unit disk $\{|t| < 1\}$, and the convergence is uniform on every disk $\{|t| < r\}$ where $r < 1$. Hence by Lemma 4.2,

$$\det \Delta_m = H(t_1, \dots, t_m) \prod_{1 \leq i < j \leq m} (t_i - t_j)^8 \tag{4.14}$$

where $H(t_1, \dots, t_m)$ is holomorphic in the unit disk.

Now, let us consider the expression of $\det \Delta_m$ in terms of the matrix elements δ_{ij} of Δ_m , that is

$$\det \Delta_m = \sum_{\sigma} \varepsilon_{\sigma} \delta_{1\sigma(1)} \cdots \delta_{2m\sigma(2m)} \tag{4.15}$$

where σ is a permutation of $\{1, \dots, 2m\}$ and $\varepsilon_{\sigma} = \pm 1$ depending on whether σ is even or odd. The common denominator of the sum in (4.15) is

$$\prod_{i=1}^m (1 - t_i^2)^4 \prod_{1 \leq i < j \leq m} (1 - t_i t_j)^8 \tag{4.16}$$

Therefore by (4.14),

$$\det \Delta_m = \frac{\prod_{1 \leq i < j \leq m} (t_i - t_j)^8 C(t_1, \dots, t_m)}{\prod_{i=1}^m (1 - t_i^2)^4 \prod_{1 \leq i < j \leq m} (1 - t_i t_j)^8} \tag{4.17}$$

where $C(t_1, \dots, t_m)$ is a polynomial of t_1, \dots, t_m . Observe that (4.17) holds for all points t_1, \dots, t_m in the unit disk, and so it can be extended to the whole complex plane. We are going to show that $C(t_1, \dots, t_m)$ is a constant, and moreover, that

$$C(t_1, \dots, t_m) = 1 \tag{4.18}$$

Let us look at the asymptotic behavior of $\det \Delta_m$ as $t_1 \rightarrow \infty$ while t_2, \dots, t_m are fixed. To see it more clearly, let us change Δ_m to $\Delta_m^{(1)}$ by subtracting the

(2*i* - 1)th column and row from (2*i*)th column and row, respectively, for *i* = 1, ..., *m*. Then

$$\Delta_m^{(1)} = \begin{pmatrix} \Delta_{11}^{(1)} & \Delta_{12}^{(1)} & \dots & \Delta_{1m}^{(1)} \\ \Delta_{21}^{(1)} & \Delta_{22}^{(1)} & \dots & \Delta_{2m}^{(1)} \\ \dots & \dots & \dots & \dots \\ \Delta_{m1}^{(1)} & \Delta_{m2}^{(1)} & \dots & \Delta_{mm}^{(1)} \end{pmatrix} \tag{4.19}$$

where

$$\Delta_{kk}^{(1)} = \begin{pmatrix} \frac{1}{1-t_k^2} & 0 \\ 0 & \frac{1}{(1-t_k^2)^3} \end{pmatrix} \tag{4.20}$$

for *k* = 1, ..., *m*, and

$$\Delta_{ij}^{(1)} = \begin{pmatrix} \frac{1}{1-t_it_j} & \frac{t_i-t_j}{(1-t_it_j)^2(1-t_j^2)} \\ \frac{t_i-t_j}{(1-t_it_j)^2(1-t_j^2)} & \frac{2t_it_j+t_i^2t_j^2-2t_j^2-2t_i^2+1}{(1-t_j^2)(1-t_i^2)(1-t_it_j)^3} \end{pmatrix} \tag{4.21}$$

for *i* ≠ *j*. The leading powers of the elements of Δ_{*m*}⁽¹⁾, as *t*₁ → ∞, are

$$\Delta_m^{(2)} = \begin{pmatrix} 1/t_1^2 & 0 & 1/t_1 & 1/t_1 & \dots & 1/t_1 & 1/t_1 \\ 0 & 1/t_1^6 & 1/t_1^3 & 1/t_1^3 & \dots & 1/t_1^3 & 1/t_1^3 \\ 1/t_1 & 1/t_1^3 & * & * & \dots & * & * \\ 1/t_1 & 1/t_1^3 & * & * & \dots & * & * \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 1/t_1 & 1/t_1^3 & * & * & \dots & * & * \\ 1/t_1 & 1/t_1^3 & * & * & \dots & * & * \end{pmatrix} \tag{4.22}$$

where *’s stand for the terms of the order of *O*(1). Therefore

$$\det \Delta_m = O\left(\frac{1}{t_1^8}\right), \quad t_1 \rightarrow \infty \tag{4.23}$$

By (4.17),

$$\det \Delta_m \sim \frac{\text{const} \cdot C(t_1, \dots, t_m)}{t_1^8}, \quad t_1 \rightarrow \infty$$

hence $C(t_1, \dots, t_m)$ is constant in t_1 . The same argument on t_2, \dots, t_m shows that it is independent of t_2, \dots, t_m , so it is indeed a constant, say C_m , i.e.,

$$\det \Delta_m = \frac{C_m \prod_{1 \leq i < j \leq m} (t_i - t_j)^8}{\prod_{i=1}^m (1 - t_i^2)^4 \prod_{1 \leq i < j \leq m} (1 - t_i t_j)^8} \tag{4.24}$$

To prove that $C_m = 1$, let us consider the asymptotic behavior of $\det \Delta_m$ as $t_1 \rightarrow 1$ with t_2, \dots, t_m fixed and close to zero. Then on the one hand, we have from (4.28) that

$$\lim_{t_1 \rightarrow 1} (1 - t_1^2)^4 \det \Delta_m = \frac{C_m}{C_{m-1}} \det \Delta_{m-1} \tag{4.25}$$

On the other hand,

$$\Delta_m^{(1)} = \begin{pmatrix} \frac{1}{(1-t_1^2)} & 0 & * & \dots & * \\ 0 & \frac{1}{(1-t_1^2)^3} & * & \dots & * \\ * & * & & & \\ \dots & \dots & & \Delta_{m-1} & \\ * & * & & & \end{pmatrix} \tag{4.26}$$

where the terms $*$ are regular at $t_1 = 1$. Hence the leading term of the Laurent series of $\det \Delta_m$ at $t_1 = 1$ is $(1 - t_1)^{-4} \det \Delta_{m-1}$, which shows that

$$\frac{C_m}{C_{m-1}} \det \Delta_{m-1} = \det \Delta_{m-1} \tag{4.27}$$

Thus $C_m = C_{m-1}$. Repeating this argument we get that

$$C_m = C_{m-1} = \dots = C_2 = C_1 = 1$$

Therefore $C_m = 1$. Proposition 4.1 is proved.

Similarly we prove the following proposition.

Proposition 4.3.

$$\Omega_m = \begin{pmatrix} \omega_{11} & \omega_{12} & \dots & \omega_{1m} \\ \omega_{21} & \omega_{22} & \dots & \omega_{2m} \\ \dots & \dots & \dots & \dots \\ \omega_{m1} & \omega_{m2} & \dots & \omega_{mm} \end{pmatrix} \tag{4.28}$$

with

$$\omega_{ii} = \frac{(1 - t_i^2)^3 \prod_{i \neq j} (1 - t_i t_j)^4}{\prod_{j \neq i} (t_i - t_j)^4} \tag{4.29}$$

and

$$\omega_{ij} = \frac{(1 - t_i^2)^2 (1 - t_j^2)^2 \prod_{k \neq i} (1 - t_i t_k)^2 \prod_{k \neq j} (1 - t_j t_k)^2}{(1 - t_k t_j) \prod_{k \neq i} (t_i - t_k)^2 \prod_{k \neq j} (t_j - t_k)^2}, \quad i \neq j \tag{4.30}$$

Put now

$$t_i = \tanh(\pi s_i), \quad i = 1, \dots, m \tag{4.31}$$

Then by (4.29) and (4.30),

$$\omega_{ii} = (1 - t_i^2)^3 \prod_{j \neq i} \coth^4 \pi(s_i - s_j) \tag{4.32}$$

$$\omega_{ij} = (1 - t_i^2)^{3/2} (1 - t_j^2)^{3/2} \frac{\prod_{k \neq i} \cosh^2 \pi(s_i - s_k) \prod_{k \neq j} \cosh^4 \pi(s_j - s_k)}{\cosh \pi(s_i - s_j)}$$

for $i \neq j$. In addition,

$$\det A_m = \frac{\prod_{1 \leq i < j \leq m} \tanh^8 \pi(s_i - s_j)}{\prod_{i=1}^m (1 - t_i^2)^4} \tag{4.33}$$

By (2.14),

$$k_m(s_1, \dots, s_m) = \frac{K_m(t_1, \dots, t_m)}{\prod_{i=1}^m p(t_i)} \tag{4.34}$$

Now let us substitute y_i 's in (4.6) by

$$(1 - t_i^2)^{3/2} y_i, \quad i = 1, \dots, m \tag{4.35}$$

then

$$K_m(t_1, \dots, t_m) = \frac{\prod_{i=1}^m p(t_i)}{2^m \prod_{1 \leq i < j \leq m} \tanh^4 \pi(s_i - s_j)} \times \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| e^{-1/2(Y\Sigma_m, Y)} dy_1 \dots dy_m \tag{4.36}$$

where

$$\Sigma_m = \begin{pmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1m} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2m} \\ \cdots & \cdots & \cdots & \cdots \\ \sigma_{m1} & \sigma_{m2} & \cdots & \sigma_{mm} \end{pmatrix} \quad (4.37)$$

with

$$\begin{aligned} \sigma_{ii} &= \prod_{j \neq i} \coth^4 \pi(s_i - s_j) \\ \sigma_{ij} &= \frac{\prod_{k \neq i} \coth^2 \pi(s_i - s_k) \prod_{k \neq j} \coth^2 \pi(s_j - s_k)}{\cosh \pi(s_i - s_j)} \end{aligned} \quad (4.38)$$

Now substitute y_i 's in the formula (4.36) by

$$y_i \prod_{j \neq i} \coth^2 \pi(s_i - s_j), \quad i = 1, \dots, m \quad (4.39)$$

then

$$\begin{aligned} K_m(t_1, \dots, t_m) &= 2^{-m} \prod_{i=1}^m p(t_i) \prod_{1 \leq i < j \leq m} \tanh^4 \pi(s_i - s_j) \\ &\times \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} |y_1 \cdots y_m| e^{-1/2(Y\Gamma_m, Y)} dy_1 \cdots dy_m \end{aligned} \quad (4.40)$$

where

$$\Gamma_m = \left(\frac{1}{\cosh \pi(s_i - s_j)} \right)_{i,j=1}^m \quad (4.41)$$

Thus

$$\begin{aligned} k_m(s_1, \dots, s_m) &= 2^{-m} \prod_{1 \leq i < j \leq m} \tanh^4 \pi(s_i - s_j) \\ &\times \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} |y_1 \cdots y_m| e^{-1/2(Y\Gamma_m, Y)} dy_1 \cdots dy_m \end{aligned} \quad (4.42)$$

Theorem 1.2 is proved.

Proof of Theorem 1.3. Let

$$\begin{aligned} f_n(t) &= c_0 + c_1 t + \cdots + c_{n-1} t^{n-1} + c_n t^n \\ g_n(t) &= c_0 t^n + c_1 t^{n-1} + \cdots + c_{n-1} t + c_n \end{aligned} \quad (4.43)$$

Then if $\tau_k \neq 0$ is a zero of $f_n(t)$ then τ_k^{-1} is a zero of $g_n(t)$. Hence if $1 < a < b$ then

$$\xi_f(a, b) = \xi_g(a^{-1}, b^{-1}) \quad (4.44)$$

where $\xi_f(a, b)$ is the number of zeros of $f_n(t)$ in the interval (a, b) . Observe that the distribution of the vector of coefficients (c_0, \dots, c_n) coincides with the one of the vector (c_n, \dots, c_0) . Hence

$$\begin{aligned} \mathbb{E}[\xi_f(a_1, b_1) \cdots \xi_f(a_m, b_m)] &= \mathbb{E}[\xi_g(a_1^{-1}, b_1^{-1}) \cdots \xi_g(a_m^{-1}, b_m^{-1})] \\ &= \mathbb{E}[\xi_f(a_1^{-1}, b_1^{-1}) \cdots \xi_f(a_m^{-1}, b_m^{-1})] \end{aligned} \quad (4.45)$$

Take $a_j = t_j$ and $b_j = t_j + \Delta t_j$, $j = 1, \dots, m$, and get $\Delta t_j \rightarrow 0$. Since

$$|a^{-1} - b^{-1}| = a^{-2} |a - b| + O(|a - b|^2), \quad a \rightarrow b$$

we deduce that

$$\frac{K_{nm}(t_1^{-1}, \dots, t_m^{-1})}{t_1^2 \cdots t_m^2} = K_{nm}(t_1, \dots, t_m) \quad (4.46)$$

Hence

$$\frac{K_{nm}(t_1^{-1}, \dots, t_m^{-1})}{p(t_1^{-1}) \cdots p(t_m^{-1})} = \frac{K_{nm}(t_1, \dots, t_m)}{p(t_1) \cdots p(t_m)} \quad (4.47)$$

because

$$p(t^{-1}) = \frac{1}{\pi |1 - t^{-2}|} = t^2 p(t)$$

This proves that

$$k_{nm}^{\text{out}}(s_1, \dots, s_m) = k_{nm}(s_1, \dots, s_m)$$

Theorem 1.3 is proved.

Proof of Theorem 1.4. Let $|t_1|, \dots, |t_l| < 1$ and $|t_{l+1}|, \dots, |t_m| > 1$. Denote by $D_{nm}(x_1, y_1, \dots, x_m, y_m; t_1, \dots, t_m)$ the joint distribution density of the vector

$$(f_n(t_1), f_n(t_1), \dots, f_n(t_l), f'_n(t_l), g_n(t_{l+1}), g'_n(t_{l+1}), \dots, g_n(t_m), g'_n(t_m))$$

Then

$$K_{nm}(t_1, \dots, t_m) = \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| \times D_{nm}(0, y_1, \dots, 0, y_m; t_1, \dots, t_l, t_{l+1}^{-1}, \dots, t_m^{-1}) dy_1 \dots dy_m \quad (4.48)$$

The covariance

$$E f_n(t_i) g_n(t_j^{-1}) = t_i^n + t_i^{n-1} t_j^{-1} + \dots + t_i t_j^{1-n} + t_j^{-n}, \quad 1 \leq i \leq l < j \leq m$$

goes to 0 as $n \rightarrow \infty$, together with the partial derivatives in t_i, t_j , while

$$\lim_{n \rightarrow \infty} E f_n(t_i) f_n(t_j) = \frac{1}{1 - t_i t_j}, \quad 1 \leq i, j \leq l \quad (4.49)$$

$$\lim_{n \rightarrow \infty} E g_n(t_i^{-1}) g_n(t_j^{-1}) = \frac{1}{1 - t_i^{-1} t_j^{-1}}, \quad l+1 \leq i, j \leq m$$

This proves that the limiting Gaussian kernel $D_m = \lim_{n \rightarrow \infty} D_{nm}$ is factored,

$$\begin{aligned} D_m(x_1, y_1, \dots, x_m, y_m; t_1, \dots, t_l, t_{l+1}^{-1}, \dots, t_m^{-1}) \\ = D_l(f)(x_1, y_1, \dots, x_l, y_l; t_1, \dots, t_l) \\ \times D_{m-l}(g)(x_{l+1}, y_{l+1}, \dots, x_m, y_m; t_{l+1}^{-1}, \dots, t_m^{-1}) \end{aligned}$$

Hence the correlation function K_m is also factored. Theorem 1.4 is proved.

5. CORRELATIONS BETWEEN ZEROS OF THE $O(1)$ RANDOM POLYNOMIAL

In the following two sections we discuss the correlation functions of real zeros and the variance of the number of real zeros of the $O(1)$ random polynomial. Let $f_n(t)$ be a $O(1)$ random polynomial, that is

$$f_n(t) = \sum_{k=0}^n c_k t^k \quad (5.1)$$

where $\{c_k\}$ are real independent Gaussian random variables with

$$E c_k = 0, \quad E c_k^2 = \sigma_k^2 = \binom{n}{k} \tag{5.2}$$

In this case (1.4) reduces to

$$\begin{aligned} A_n(t) &= (1 + t^2)^n \\ B_n(t) &= nt(1 + t^2)^{n-1} \\ C_n(t) &= n(1 + nt^2)(1 + t^2)^{n-2} \end{aligned} \tag{5.3}$$

which gives

$$p_n(t) = \frac{\sqrt{n}}{\pi(1 + t^2)} \tag{5.4}$$

(see [BS], [EK], [BBL1], and others). The average value of real zeros is

$$E \# \{k: \tau_k \in \mathbb{R}\} = \int_{-\infty}^{\infty} p_n(t) dt = \sqrt{n} \tag{5.5}$$

and the normalized density,

$$\frac{p_n(t)}{\sqrt{n}} = \frac{1}{\pi(1 + t^2)} \tag{5.6}$$

is the Cauchy distribution density. Observe that both (5.4) and (5.5) are exact relations for all n . Let us compute the two-point correlation function $K_{n2}(t_1, t_2)$.

Define

$$\varphi_n(t) = \frac{f_n(t)}{(1 + t^2)^{n/2}} \tag{5.7}$$

Then the real zeros of $\varphi_n(t)$ coincide with those of $f_n(t)$. Hence, similar to (2.6), we can write that

$$K_{n2}(t_1, t_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| D_{n2}(0, y_1, 0, y_2; t_1, t_2) dy_1 dy_2 \tag{5.8}$$

where $D_{n2}(x_1, y_1, x_2, y_2; t_1, t_2)$ is the distribution density function of the vector

$$(\varphi_n(t_1), \varphi'_n(t_1), \varphi_n(t_2), \varphi'_n(t_2))$$

Observe that

$$E \varphi_n(t_1) \varphi_n(t_2) = \rho_n(t_1, t_2) \equiv \frac{(1 + t_1 t_2)^n}{(1 + t_1^2)^{n/2} (1 + t_2^2)^{n/2}} \tag{5.9}$$

and

$$\begin{aligned} E \varphi'_n(t_1) \varphi_n(t_2) &= \frac{\partial \rho_n}{\partial t_1}(t_1, t_2) = n \rho_n \frac{(t_2 - t_1)}{(1 + t_1 t_2)(1 + t_1^2)} \\ E \varphi'_n(t_1) \varphi'_n(t_2) &= \frac{\partial^2 \rho_n}{\partial t_1 \partial t_2}(t_1, t_2) \\ &= -n^2 \rho_n \frac{(t_2 - t_1)^2}{(1 + t_1 t_2)^2 (1 + t_1^2)(1 + t_2^2)} + n \rho_n \frac{1}{(1 + t_1 t_2)^2} \end{aligned} \tag{5.10}$$

Define the random variable $\psi_n(t)$ as

$$\psi_n(t) = \frac{\varphi'_n(t)(1 + t^2)}{\sqrt{n}} \tag{5.11}$$

Let $D_{n2}(x_1, y_1, x_2, y_2; t_1, t_2)$ be the distribution density function of the random vector $(\varphi_n(t_1), \psi_n(t_1), \varphi_n(t_2), \psi_n(t_2))$. Then after a change of variables, (5.8) is rewritten as

$$\frac{K_{n2}(t_1, t_2)}{\rho_n(t_1) \rho_n(t_2)} = \pi^2 \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| D_{n2}(0, y_1, 0, y_2; t_1, t_2) dy_1 dy_2 \tag{5.12}$$

From (5.10),

$$\begin{aligned} E \psi_n(t_1) \varphi_n(t_2) &= \rho_n \frac{\sqrt{n}(t_2 - t_1)}{(1 + t_1 t_2)} \\ E \psi_n(t_1) \psi_n(t_2) &= \rho_n \left[-\frac{n(t_2 - t_1)^2}{(1 + t_1 t_2)^2} + \frac{(1 + t_1^2)(1 + t_2^2)}{(1 + t_1 t_2)^2} \right] \end{aligned} \tag{5.13}$$

hence the covariance matrix of the vector $(\varphi_n(t_1), \psi_n(t_1), \varphi_n(t_2), \psi_n(t_2))$ is

$$\Delta_n = \begin{pmatrix} 1 & 0 & \rho_n & \rho_n \frac{\sqrt{n}(t_1 - t_2)}{(1 + t_1 t_2)} \\ 0 & 1 & \rho_n \frac{\sqrt{n}(t_2 - t_1)}{(1 + t_1 t_2)} & \rho_n a \\ \rho_n & \rho_n \frac{\sqrt{n}(t_2 - t_1)}{(1 + t_1 t_2)} & 1 & 0 \\ \rho_n \frac{\sqrt{n}(t_1 - t_2)}{(1 + t_1 t_2)} & \rho_n a & 0 & 1 \end{pmatrix} \quad (5.14)$$

where

$$a = a(t_1, t_2) = -\frac{n(t_2 - t_1)^2}{(1 + t_1 t_2)^2} + \frac{(1 + t_1^2)(1 + t_2^2)}{(1 + t_1 t_2)^2} \quad (5.15)$$

Suppose that t_1 and t_2 are two distinct fixed points. Then as $n \rightarrow \infty$, the quantity

$$|\rho_n| = \left[1 - \frac{(t_1 - t_2)^2}{(1 + t_1^2)(1 + t_2^2)} \right]^{n/2} \quad (5.16)$$

goes to 0 exponentially fast, and hence Δ_n approaches the unit matrix exponentially fast. By (5.12) this implies that

$$\lim_{n \rightarrow \infty} \frac{K_{n2}(t_1, t_2)}{p_n(t_1) p_n(t_2)} = 1$$

and the rate of convergence is exponential. In the same way we obtain that if t_1, \dots, t_m are m distinct fixed points then

$$\begin{aligned} & \frac{K_{nm}(t_1, \dots, t_m)}{p_n(t_1) \cdots p_n(t_m)} \\ &= \pi^m \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} |y_1 \cdots y_m| D_{nm}(0, y_1, \dots, 0, y_m; t_1, \dots, t_m) dy_1 \cdots dy_m \end{aligned} \quad (5.17)$$

where $D_{nm}(x_1, y_1, \dots, x_m, y_m; t_1, \dots, t_m)$ is a $(2m) \times (2m)$ Gaussian density with the covariance matrix

$$A_n = (A_{ij}^{(n)})_{i,j=1,\dots,m}$$

where

$$A_{ii}^{(n)} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A_{ij}^{(n)} = \rho_n(t_i, t_j) \begin{pmatrix} 1 & \frac{\sqrt{n}(t_i - t_j)}{(1 + t_i t_j)} \\ \frac{\sqrt{n}(t_j - t_i)}{(1 + t_i t_j)} & a(t_i, t_j) \end{pmatrix} \quad (5.18)$$

where $\rho_n(t_1, t_2)$ and $a(t_1, t_2)$ are defined in (5.9) and (5.15), respectively. For fixed different t_1, \dots, t_m the matrix A_n approaches the unit matrix exponentially fast, and this implies that

$$\lim_{n \rightarrow \infty} \frac{K_{nm}(t_1, \dots, t_m)}{p_n(t_1) \cdots p_n(t_m)} = 1$$

and the rate of convergence is exponential. This means the independence of the distribution of real zeros at distinct fixed points.

The formula (5.17) is simplified if we make the change of variables $t = \tan \theta$ (stereographic projection). Consider therefore the random function

$$g_n(\theta) = \sum_{j=0}^n c_j \sin^j(\theta) \cos^{n-j}(\theta), \quad -\frac{\pi}{2} < \theta \leq \frac{\pi}{2} \quad (5.19)$$

Then

$$g_n(\theta) = \cos^n \theta f_n(\tan \theta) \quad (5.20)$$

hence if $\{\tau_j\}$ are zeros of $f_n(t)$ then

$$\{\eta_j = \arctan \tau_j\} \quad (5.21)$$

are zeros of $g_n(\theta)$. When $c_n = 0$, $g_n(\theta)$ has an extra zero $\eta = \pi/2$. Since the probability that $c_n = 0$ is equal to zero, the joint probability distributions of zeros of the functions $g_n(\theta)$ and $f_n(\tan \theta)$ coincide. By (5.6) the zeros $\{\eta_j\}$ are uniformly distributed on the interval $[-\pi/2, \pi/2]$. The function $g_n(\theta)$ is a Gaussian random function with zero mean and the covariance function

$$\begin{aligned} E g_n(\theta_1) g_n(\theta_2) &= \sum_{k=0}^n \binom{n}{k} \sin^k \theta_1 \cos^{n-k} \theta_1 \sin^k \theta_2 \cos^{n-k} \theta_2 \\ &= \cos^n(\theta_1 - \theta_2) \end{aligned} \quad (5.22)$$

The function $g_n(\theta)$ is periodic of period π if n is even, and it is periodic of period 2π if n is odd. It is convenient to consider $g_n(\theta)$ on the circle $S^1 = \mathbb{R}/(2\pi)\mathbb{Z}$ of the length 2π . This circle is the covering space of the original circle $\mathbb{R}/\pi\mathbb{Z}$. If $g_n(\theta) = 0$ then $g_n(\theta + \pi) = 0$ as well. On S^1 the distribution of $g_n(\theta)$ is invariant with respect to the shift

$$\theta \rightarrow \alpha + \theta \pmod{2\pi}$$

hence it is $O(1)$ -invariant.

Let $K_{nm}(\theta_1, \dots, \theta_m)$ be the correlation function of the zeros $\{\eta_j\}$ of $g_n(\theta)$. Assume that

$$\theta_j - \theta_k \neq 0 \pmod{\pi}, \quad 1 \leq j, k \leq m \tag{5.23}$$

Then (5.17) gives that

$$\begin{aligned} & \frac{K_{nm}(\theta_1, \dots, \theta_m)}{(\pi^{-1} \sqrt{n})^m} \\ &= \pi^m \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| D_{nm}(0, y_1, \dots, 0, y_m; \theta_1, \dots, \theta_m) dy_1 \dots dy_m \end{aligned} \tag{5.24}$$

where $D_{nm}(x_1, y_1, \dots, x_m, y_m; \theta_1, \dots, \theta_m)$ is a $(2m) \times (2m)$ Gaussian density with the covariance matrix

$$A_n = (A_{ij}^{(n)})_{i,j=1,\dots,m}$$

where

$$A_{ij}^{(n)} = \cos^n(\theta_i - \theta_j) \begin{pmatrix} 1 & \sqrt{n} \tan(\theta_i - \theta_j) \\ \sqrt{n} \tan(\theta_j - \theta_i) & -n \tan^2(\theta_i - \theta_j) + \cos^{-2}(\theta_i - \theta_j) \end{pmatrix} \tag{5.25}$$

Observe that $D_{nm}(x_1, y_1, \dots, x_m, y_m; \theta_1, \dots, \theta_m)$ is the probability distribution density of the vector

$$\left(g_n(\theta_1), \frac{g'_n(\theta_1)}{\sqrt{n}}, \dots, g_n(\theta_m), \frac{g'_n(\theta_m)}{\sqrt{n}} \right)$$

and it is nondegenerate provided that $n \geq 2m - 1$ and (5.23) holds.

Consider now the scaling limit of the correlation functions. The straightened zeros are

$$\zeta_j = \frac{\eta_j \sqrt{n}}{\pi}$$

They are uniformly distributed on the circle of the length $2\sqrt{n}$. The limit m -point correlation function of $\{\zeta_j\}$ is

$$k_m(s_1, \dots, s_m) = \lim_{n \rightarrow \infty} \frac{K_{nm}(\theta_1, \dots, \theta_m)}{(\pi^{-1} \sqrt{n})^m}, \quad \theta_i = \frac{s_i \pi}{\sqrt{n}} \tag{5.26}$$

Let us find $k_m(s_1, \dots, s_m)$. We have that

$$\lim_{n \rightarrow \infty} \cos^n \left(\frac{(s_i - s_j) \pi}{\sqrt{n}} \right) = e^{-\pi^2 (s_i - s_j)^2 / 2} \tag{5.27}$$

and by (5.25),

$$\lim_{n \rightarrow \infty} \Delta_n = \Delta = (\Delta_{ij})_{i,j=1,\dots,m}$$

with

$$\Delta_{ij} = e^{-\pi^2 (s_i - s_j)^2 / 2} \begin{pmatrix} 1 & \pi(s_i - s_j) \\ \pi(s_j - s_i) & 1 - \pi^2 (s_i - s_j)^2 \end{pmatrix} \tag{5.28}$$

By (5.24) this gives that

$$k_m(s_1, \dots, s_m) = \pi^m \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| d_m(0, y_1, \dots, 0, y_m; s_1, \dots, s_m) dy_1 \dots dy_m \tag{5.29}$$

and $d_m(x_1, y_1, \dots, x_m, y_m; s_1, \dots, s_m)$ is a Gaussian density with the covariance matrix Δ . Let Ω be $m \times m$ matrix obtained by deleting all odd rows and odd columns from the matrix Δ^{-1} . Then we can write (5.29) as

$$k_m(s_1, \dots, s_m) = \frac{1}{2^m (\det \Delta)^{1/2}} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} |y_1 \dots y_m| e^{-1/2 (Y\Omega, Y)} dy_1 \dots dy_m \tag{5.30}$$

In Appendix C below we prove that

$$\Delta > 0 \quad \text{if } s_i \neq s_j, \quad 1 \leq i < j \leq m \quad (5.31)$$

so the formula (5.30) is well-defined when the point s_i are distinct.

For $m = 2$, (5.30) reduces to

$$k_2(s_1, s_2) = \frac{1}{4\pi^2(\det \Delta)^{1/2}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(Y\Omega, Y)} dy_1 dy_2 \quad (5.32)$$

where

$$\Delta = \begin{pmatrix} 1 & 0 & e^{-s^2/2} & se^{-s^2/2} \\ 0 & 1 & -se^{-s^2/2} & (1-s^2)e^{-s^2/2} \\ e^{-s^2/2} & -se^{-s^2/2} & 1 & 0 \\ se^{-s^2/2} & (1-s^2)e^{-s^2/2} & 0 & 1 \end{pmatrix} \quad (5.33)$$

$$s = \pi(s_1 - s_2)$$

and

$$\Omega = (\Delta^{-1})_{\{2,4\}}$$

i.e., Ω is the 2×2 -submatrix of Δ^{-1} at the second and the fourth rows and columns. Observe that

$$\det \Omega = \frac{\det \Delta_{\{1,3\}}}{\det \Delta} = \frac{1 - e^{-\pi^2 s^2}}{\det \Delta}$$

A direct computation gives

$$\det \Delta = (1 - e^{-s^2})^2 - s^4 e^{-s^2} \quad (5.34)$$

and

$$\Omega = \begin{pmatrix} A & B \\ B & A \end{pmatrix}$$

where

$$A = \frac{1 - e^{-s^2} - s^2 e^{-s^2}}{\det \Delta}, \quad B = \frac{e^{-s^2/2}(e^{-s^2} + s^2 - 1)}{\det \Delta} \quad (5.35)$$

Since

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(Ay_1^2 + Ay_2^2 + 2By_1 y_2)} dy_1 dy_2 = \frac{4}{\det \Omega} \left(1 + \frac{\delta}{\sqrt{1-\delta^2}} \arcsin \delta \right)$$

where $\delta = B/A$ (see Appendix A), we obtain from (5.26) that

$$\begin{aligned} k_2(s_1, s_2) &= \frac{1}{(\det A)^{1/2} \det \Omega} \left(1 + \frac{\delta}{\sqrt{1-\delta^2}} \arcsin \delta \right) \\ &= \frac{(\det A)^{1/2}}{1 - e^{-s^2}} \left(1 + \frac{\delta}{\sqrt{1-\delta^2}} \arcsin \delta \right) \\ &= \frac{[(1 - e^{-s^2})^2 - s^4 e^{-s^2}]^{1/2}}{1 - e^{-s^2}} \left(1 + \frac{\delta}{\sqrt{1-\delta^2}} \arcsin \delta \right) \end{aligned} \tag{5.36}$$

where $s = \pi(s_1 - s_2)$ and

$$\delta = \frac{e^{-s^2/2}(e^{-s^2} + s^2 - 1)}{1 - e^{-s^2} - s^2 e^{-s^2}} \tag{5.37}$$

It is worth to note that Hannay [Han] has calculated the limit two-point correlation function of zeros for the complex random $SU(2)$ polynomial, and our calculation of (5.36) is very similar to the one of Hannay.

As $s \rightarrow 0$,

$$\det A = \frac{s^8}{12} + O(s^{10}), \quad \delta = 1 - \frac{s^2}{6} + O(s^4)$$

which implies that

$$k_2(s_1, s_2) = \frac{\pi^2 |s_1 - s_2|}{4} + O(|s_1 - s_2|^2), \quad s_1 - s_2 \rightarrow 0 \tag{5.38}$$

As $s \rightarrow \infty$,

$$\det A = 1 - s^4 e^{-s^2} + O(e^{-s^2}), \quad \delta = -s^2 e^{-s^2/2} + O(e^{-s^2/2})$$

which implies that

$$k_2(s_1, s_2) = 1 + \frac{\pi^4(s_1 - s_2)^4 e^{-\pi^2(s_1 - s_2)^2}}{2} + O((s_1 - s_2)^2 e^{-\pi^2(s_1 - s_2)^2}) \tag{5.39}$$

$$|s_1 - s_2| \rightarrow \infty$$

Thus we have proved the following theorem.

Theorem 5.1. Let $\{\tau_j\}$ be zeros of a random $O(1)$ polynomial $f_n(t)$ of degree n , and

$$\zeta_j = \frac{\sqrt{n} \arctan \tau_j}{\pi}$$

be the straightened zeros. Then the limit m -point correlation function $k_m(s_1, \dots, s_m)$ of $\{\zeta_j\}$ is given by the formula (5.30) where Δ is a $(2m) \times (2m)$ symmetric matrix which consists of 2×2 blocks Δ_{ij} defined in (5.27), and $\Omega = (\Delta^{-1})_{\text{even}}$ is the $m \times m$ matrix of the elements of Δ^{-1} with even indices. The 2-point correlation function is given by the formula (5.36), and its asymptotics as $s_1 - s_2 \rightarrow 0$ and $|s_1 - s_2| \rightarrow \infty$ are given in (5.38) and (5.39), respectively.

The graph of $k_2(0, s)$ is shown in Fig. 2.




Fig. 2. The two-point correlation function of straightened zeros of the $O(1)$ random polynomial.

6. VARIANCE OF THE NUMBER OF REAL ZEROS OF THE $O(1)$ RANDOM POLYNOMIAL

Here we calculate the variance of the random variable $\xi_n(a, b)$ in the case when $f_n(t)$ is the $O(1)$ random polynomial. By definition,

$$\text{Var } \xi_n(a, b) = E \xi_n^2(a, b) - (E \xi_n(a, b))^2 \tag{6.1}$$

By (2.17),

$$E \xi_n^2(a, b) = \int_a^b p_n(t) dt + \int_a^b \int_a^b K_{n2}(t_1, t_2) dt_1 dt_2 \tag{6.2}$$

Since

$$(E \xi_n(a, b))^2 = \int_a^b p_n(t_1) dt_1 \int_a^b p_n(t_2) dt_2 \tag{6.3}$$

we obtain that

$$\begin{aligned} \text{Var } \xi_n(a, b) &= \int_a^b p_n(t) dt + \int_a^b \int_a^b [K_{n2}(t_1, t_2) - p_n(t_1) p_n(t_2)] dt_1 dt_2 \\ &= \int_a^b p_n(t) dt + \int_a^b \int_a^b \left[\frac{K_{n2}(t_1, t_2)}{p_n(t_1) p_n(t_2)} - 1 \right] p_n(t_1) p_n(t_2) dt_1 dt_2 \end{aligned} \tag{6.4}$$

When t_1 and t_2 are separated, the difference

$$\frac{K_{n2}(t_1, t_2)}{p_n(t_1) p_n(t_2)} - 1$$

is exponentially small, hence the main contribution to the last integral comes from close t_1, t_2 . Let us put

$$t_1 = t, \quad t_2 = t + \frac{s}{p_n(t)}$$

Then

$$\begin{aligned} &\int_a^b \int_a^b \left[\frac{K_{n2}(t_1, t_2)}{p_n(t_1) p_n(t_2)} - 1 \right] p_n(t_1) p_n(t_2) dt_1 dt_2 \\ &\sim \int_a^b p_n(t) dt \int_{-\infty}^{\infty} (k_2(0, s) - 1) ds \end{aligned}$$

hence

$$\text{Var } \xi_n(a, b) \sim \int_a^b p_n(t) dt \left[1 - \int_{-\infty}^{\infty} (1 - k_2(0, s)) ds \right]$$

Thus

$$\text{Var } \xi_n(a, b) \sim C \sqrt{n} \int_a^b \frac{dt}{\pi(1+t^2)}$$

where

$$C = 1 - \int_{-\infty}^{\infty} (k_2(0, s) - 1) ds$$

and $k_2(s_1, s_2)$ is the two-point correlation function given in (5.35). In particular,

$$\text{Var } \xi_n(-\infty, \infty) \sim C \sqrt{n}$$

Numerical value of C is

$$C = 0.5717310486902\dots$$

APPENDIX A. CALCULATION OF AN INTEGRAL

In this Appendix we show that

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(Ay_1^2 + Cy_2^2 + 2By_1 y_2)} dy_1 dy_2 \\ &= \frac{4}{AC - B^2} \left(1 + \frac{\delta}{\sqrt{1 - \delta^2}} \arcsin \delta \right) \end{aligned} \quad (\text{A.1})$$

where $\delta = B/\sqrt{AC}$. By a change of variables we can reduce the integral (A.1) to

$$I = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y_1 y_2| e^{-1/2(y_1^2 + y_2^2 + 2\delta y_1 y_2)} dy_1 dy_2$$

Changing then

$$y_1 = \frac{1}{\sqrt{2}}(x_1 + x_2)$$

$$y_2 = \frac{1}{\sqrt{2}}(x_1 - x_2)$$

we obtain that

$$I = \frac{1}{2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |x_1^2 - x_2^2| e^{-1/2[(1+\delta)x_1^2 + (1-\delta)x_2^2]} dx_1 dx_2$$

Let

$$x_1 = \frac{r}{\sqrt{1+\delta}} \cos \theta$$

$$x_2 = \frac{r}{\sqrt{1-\delta}} \sin \theta$$

Then

$$I = \frac{1}{2(1-\delta^2)^{1/2}} \int_0^{\infty} r^3 e^{-r^2/2} dr \int_0^{2\pi} \left| \frac{\cos^2 \theta}{1+\delta} - \frac{\sin^2 \theta}{1-\delta} \right| d\theta$$

$$= \frac{1}{(1-\delta^2)^{3/2}} \int_0^{2\pi} |(1-\delta) \cos^2 \theta - (1+\delta) \sin^2 \theta| d\theta$$

$$= \frac{1}{(1-\delta^2)^{3/2}} \int_0^{2\pi} |\delta - \cos 2\theta| d\theta$$

Evaluating the last integral we obtain that

$$I = \frac{4}{(1-\delta^2)^{3/2}} (\sqrt{1-\delta^2} + \delta \arcsin \delta)$$

hence (A.1) is proved.

APPENDIX B. POSITIVITY OF THE COVARIANCE MATRIX

Assume that $f_n(t) = c_0 + c_1 t + \dots + c_n t^n$ is a polynomial with independent random coefficients such that $E c_k = 0$ and $0 < E c_k^2 < \infty$. We show in this Appendix that the covariance matrix of the random vector

$$F_n = (f_n(t_1), f'_n(t_1), \dots, f_n(t_m), f'_n(t_m)) \quad (\text{B.1})$$

is positive, if t_1, \dots, t_m are pairwise different and $n \geq 2m - 1$. Consider the real-valued quadratic form associated with the covariance matrix,

$$\begin{aligned} Q(\alpha, \beta) &= \sum_{j,k} [\alpha_j \alpha_k \mathbb{E} f_n(t_j) f_n(t_k) + \alpha_j \beta_k \mathbb{E} f_n(t_j) f'_n(t_k) \\ &\quad + \beta_j \alpha_k \mathbb{E} f'_n(t_j) f_n(t_k) + \beta_j \beta_k \mathbb{E} f'_n(t_j) f'_n(t_k)] \\ &= \mathbb{E} \left| \sum_j [\alpha_j f_n(t_j) + \beta_j f'_n(t_j)] \right|^2 \\ &= \sum_{k=0}^n \sigma_k^2 \left[\sum_{j=1}^m (\alpha_j t_j^k + \beta_j k t_j^{k-1}) \right]^2 \end{aligned}$$

The generalized Vandermonde matrix

$$(t_j^k, k t_j^{k-1})_{j=1, \dots, m; k=0, \dots, 2m-1}$$

is nondegenerate, hence

$$\sum_{k=0}^n \sigma_k^2 \left[\sum_{j=1}^m (\alpha_j t_j^k + \beta_j k t_j^{k-1}) \right]^2 > 0$$

provided that not all α_j, β_j are zero. This proves that the quadratic form $Q(\alpha, \beta)$ is positive, hence the covariance matrix of the random vector (B.1) is positive as well.

The proof remains valid if c_k are, in general, dependent random variables with positive covariance matrix $(V_{kl} = \mathbb{E} c_k c_l)_{k,l=0, \dots, n}$. In this case

$$Q(\alpha, \beta) = \sum_{k,l=0}^n V_{kl} d_k d_l > 0, \quad d_k = \sum_{j=1}^m (\alpha_j t_j^k + \beta_j k t_j^{k-1})$$

provided that $n \geq 2m - 1$ and not all α_j, β_j are zero.

APPENDIX C. PROOF OF THE INEQUALITY $\Delta > 0$

Let $\Delta = (\Delta_{jk})_{j,k=1, \dots, m}$ where

$$\Delta_{jk} = e^{-(s_j - s_k)^2/2} \begin{pmatrix} 1 & (s_j - s_k) \\ (s_k - s_j) & 1 - (s_j - s_k)^2 \end{pmatrix}$$

We prove in this Appendix that $\Delta > 0$. Consider the complex-valued quadratic form associated with Δ ,

$$Q(\alpha, \beta) = \sum_{j,k} e^{-(s_j - s_k)^2/2} [\alpha_j \bar{\alpha}_k + (s_j - s_k) \alpha_j \bar{\beta}_k + (s_k - s_j) \beta_j \bar{\alpha}_k + [1 - (s_j - s_k)^2] \beta_j \bar{\beta}_k]$$

Using the formulae

$$\begin{aligned} e^{-(s_j - s_k)^2/2} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i(s_j - s_k)x} e^{-x^2/2} dx \\ (s_j - s_k) e^{-(s_j - s_k)^2/2} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i(s_j - s_k)x} (-ix) e^{-x^2/2} dx \\ [1 - (s_j - s_k)^2] e^{-(s_j - s_k)^2/2} &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{i(s_j - s_k)x} x^2 e^{-x^2/2} dx \end{aligned}$$

we can rewrite Q as

$$Q(\alpha, \beta) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx e^{-x^2/2} \sum_{j,k} [\alpha_j \bar{\alpha}_k e^{i(s_j - s_k)x} + \alpha_j \bar{\beta}_k e^{i(s_j - s_k)x} (-ix) + \beta_j \bar{\alpha}_k e^{i(s_j - s_k)x} (ix) + \beta_j \bar{\beta}_k e^{i(s_j - s_k)x} x^2]$$

Define

$$f(x) = \sum_{j=1}^m \alpha_j e^{is_j x}, \quad g(x) = \sum_{j=1}^m \beta_j e^{is_j x}$$

Then

$$Q(\alpha, \beta) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} |f(x) + ixg(x)|^2 e^{-x^2/2} dx$$

The function $f(x) + ixg(x)$ is not identically zero provided that not all α_j, β_j are 0. Indeed, for every test function $\varphi(x)$,

$$\int_{-\infty}^{\infty} (f(x) + ixg(x)) \varphi(x) dx = \sum_{j=1}^m (\alpha_j \tilde{\varphi}(s_j) + \beta_j \tilde{\varphi}'(s_j))$$

where $\tilde{\varphi}(s)$ is the Fourier transform of $\varphi(x)$. We can localize the function $\tilde{\varphi}(s)$ near s_j and make the last sum nonzero. This proves that $f(x) + ixg(x)$ is not identically zero, and hence $Q(\alpha, \beta) > 0$. Hence $\Delta > 0$.

ACKNOWLEDGMENTS

This work was supported in part by the National Science Foundation, Grant No. DMS-9623214, and this support is gratefully acknowledged.

REFERENCES

- [BS] A. T. Bharucha-Reid and M. Sambandham, *Random polynomials*, Academic Press, New York, 1986.
- [BP] A. Bloch and G. Polya, On the roots of certain algebraic equations, *Proc. London Math. Soc.* (3) **33**:102–114 (1932).
- [BBL1] E. Bogomolny, O. Bohigas, and P. Leboeuf, Distribution of roots of random polynomials, *Phys. Rev. Lett.* **68**:2726–2729 (1992).
- [BBL2] E. Bogomolny, O. Bohigas, and P. Leboeuf, Quantum chaotic dynamics and random polynomials. Preprint, 1996.
- [Dys] F. J. Dyson, Correlation between eigenvalues of a random matrix, *Commun. Math. Phys.* **19**:235–250 (1970).
- [EK] A. Edelman and E. Kostlan, How many zeros of a random polynomial are real? *Bull. Amer. Math. Soc.* **32**:1–37 (1995).
- [EO] P. Erdős and A. Offord, On the number of real roots of random algebraic equation, *Proc. London Math. Soc.* **6**:139–160 (1956).
- [ET] P. Erdős and P. Turán, On the distribution of roots of polynomials, *Ann. Math.* **51**:105–119 (1950).
- [Han] J. H. Hannay, Chaotic analytic zero points: exact statistics for those of a random spin state, *J. Phys. A: Math. Gen.* **29**:101–105 (1996).
- [IM] I. A. Ibragimov and N. B. Maslova, On the average number of real zeros of random polynomials. I, II, *Teor. Veroyatn. Primen.* **16**:229–248, 495–503 (1971).
- [K1] M. Kac, On the average number of real roots of a random algebraic equation, *Bull. Amer. Math. Soc.* **49**:314–320 (1943).
- [K2] M. Kac, On the average number of real roots of a random algebraic equation. II, *Proc. London Math. Soc.* **50**:390–408 (1948).
- [K3] M. Kac, *Probability and related topics in physical sciences*. Wiley (Interscience), New York, 1959.
- [Kos] E. Kostlan, On the distribution of roots of random polynomials, *From Topology to Computation: Proceedings of the Smalefest*, M. W. Hirsh, J. E. Marsden, and M. Shub, eds. (Springer-Verlag, New York, 1993), pp. 419–431.
- [Leb1] P. Leboeuf, Phase space approach to quantum dynamics, *J. Phys. A: Math. Gen.* **24**:4575 (1991).
- [Leb2] P. Leboeuf, Statistical theory of chaotic wavefunctions: a model in terms of random analytic functions. Preprint, 1996.
- [LS] P. Leboeuf and P. Shukla, Universal fluctuations of zeros of chaotic wavefunctions. Preprint, 1996.
- [LO] J. Littlewood and A. Offord, On the number of real roots of random algebraic equation. I, II, III, *J. London Math. Soc.* **13**:288–295 (1938); *Proc. Camb. Phil. Soc.* **35**:133–148 (1939); *Math. Sborn.* **12**:277–286 (1943).
- [LS] B. F. Logan and L. A. Shepp, Real zeros of a random polynomial. I, II, *Proc. London Math. Soc.* **18**:29–35, 308–314 (1968).
- [Maj] E. Majorana, *Nuovo Cimento* **9**:43–50 (1932).

- [Mas1] N. B. Maslova, On the variance of the number of real roots of random polynomials, *Teor. Veroyatn. Primen.* **19**:36–51 (1974).
- [Mas2] N. B. Maslova, On the distribution of the number of real roots of random polynomials, *Teor. Veroyatn. Primen.* **19**:488–500 (1974).
- [Meh] M. L. Mehta, *Random matrices* (Academic Press, New York, 1991).
- [M-A] G. Mezincescu, D. Bessis, J. Fournier, G. Mantica, and F. D. Aaron, Distribution of roots of random polynomials. Preprint, 1996.
- [SS] M. Shub and S. Smale, Complexity of Bezout's Theorem II: Volumes and Probabilities, *Computational Algebraic Geometry* (F. Eysette and A. Calligo, eds.), Progr. Math., vol. 109, Birkhauser, Boston, 1993, pp. 267–285.
- [Ste] D. Stevens, Expected number of real zeros of polynomials, *Notices Amer. Math. Soc.* **13**:79 (1966).